



Computational quest for understanding the role of astrocyte signaling in synaptic transmission and plasticity

Maurizio De Pittà^{1*}, Vladislav Volman^{2,3,4}, Hugues Berry⁵, Vladimir Parpura^{6,7}, Andrea Volterra⁸ and Eshel Ben-Jacob^{1,9,10*}

¹ School of Physics and Astronomy, Tel Aviv University, Ramat Aviv, Israel

² Center for Theoretical Biological Physics, University of California at San Diego, La Jolla, CA, USA

³ Computational Neurobiology Laboratory, The Salk Institute, La Jolla, CA, USA

⁴ L3 Applied Technologies: Simulation, Engineering and Testing, San Diego, CA, USA

⁵ BEAGLE, INRIA Rhône-Alpes, Université de Lyon, LIRIS, UMR5205, Villeurbanne, France

⁶ Department of Neurobiology, University of Alabama, Birmingham, AL, USA

⁷ Department of Biotechnology, University of Rijeka, Rijeka, Croatia

⁸ Department of Basic Neurosciences, University of Lausanne, Lausanne, Switzerland

⁹ Center for Theoretical Biological Physics, Rice University, Houston, TX, USA

¹⁰ Research and Development Unit, Assaf Harofeh Medical Center, Zerifin, Israel

Edited by:

Liam J. McDaid, University of Ulster, UK

Reviewed by:

Florentin Wörgötter, University Goettingen, Germany

Vladimir Brezina, Mount Sinai School of Medicine, USA

*Correspondence:

Maurizio De Pittà and Eshel Ben-Jacob, School of Physics and Astronomy, Tel Aviv University, 69978 Ramat Aviv, Israel.
e-mail: mauriodepitta@gmail.com; eshelbj@gmail.com

The complexity of the signaling network that underlies astrocyte-synapse interactions may seem discouraging when tackled from a theoretical perspective. Computational modeling is challenged by the fact that many details remain hitherto unknown and conventional approaches to describe synaptic function are unsuitable to explain experimental observations when astrocytic signaling is taken into account. Supported by experimental evidence is the possibility that astrocytes perform genuine information processing by means of their calcium signaling and are players in the physiological setting of the basal tone of synaptic transmission. Here we consider the plausibility of this scenario from a theoretical perspective, focusing on the modulation of synaptic release probability by the astrocyte and its implications on synaptic plasticity. The analysis of the signaling pathways underlying such modulation refines our notion of tripartite synapse and has profound implications on our understanding of brain function.

Keywords: astrocyte-synapse interactions, astrocyte modeling, calcium signaling, calcium encoding, gliotransmission, synaptic plasticity, metaplasticity, cortical maps

INTRODUCTION

The simultaneous recognition that astrocytes sense neighboring neuronal activity and release neuroactive agents (or “gliotransmitters”) has been instrumental in the uncovering of the many roles played by these cells in the control of genesis, function and plasticity of synapses (Haydon, 2001; Ullian et al., 2004; Volterra and Meldolesi, 2005; Bains and Oliet, 2007; Santello and Volterra, 2009; Zorec et al., 2012). These findings initiated a conceptual revolution that leads to rethinking how brain communication works since they imply that information travels and is processed not just in the neuronal circuitry but in an expanded neuron-glia

network (Haydon, 2001; Volterra and Meldolesi, 2005; Giaume et al., 2010). On the other hand the physiological need for astrocyte signaling in brain information processing and the modes of action of these cells in computational tasks remain largely undefined. This is due, to a large extent, both to the lack of conclusive experimental evidence, and to a substantial lack of a theoretical framework to address modeling and characterization of the many possible astrocyte functions. This review aims at introducing such a perspective providing a framework for future modeling efforts in the field based on preliminary theoretical studies on both astrocytic calcium signaling and gliotransmitter-mediated modulations of synaptic release probability.

A THEORETICAL FRAMEWORK FOR ASTROCYTE-SYNAPSE INTERACTIONS

Control of synaptic transmission and plasticity by astrocytes subtends a complex signaling network, which involves different biochemical pathways (Volterra and Meldolesi, 2005; Zorec et al., 2012). In general, synaptically-released neurotransmitter can spill out of the synaptic cleft and bind to metabotropic receptors found on the neighboring astrocytic processes triggering there inositol 1,4,5-trisphosphate (IP₃)-mediated Ca²⁺ signaling. This was observed at both glutamatergic, cholinergic, noradrenergic, and GABAergic synapses in the hippocampus,

Abbreviations: Adn, adenosine; AFM, amplitude and frequency modulation; AM, amplitude modulation; AMPA, 2-amino-3-(5-methyl-3-oxo-1,2-oxazol-4-yl) propanoic acid; AMPAR, AMPA glutamate receptor; AP, action potential; ATP, adenosine triphosphate; Ca²⁺, calcium; CICR, Ca²⁺-induced Ca²⁺ release; D-ser, D-serine; EAAT, excitatory aminoacid transporter; ER, endoplasmic reticulum; FM, frequency modulation; GABA, γ -aminobutyric acid; GJC, gap-junction channel; Glu, glutamate; GluR, glutamate receptor; mGluRs, metabotropic glutamate receptor; GPCR, G_q protein-coupled receptor; IP-5P, inositol polyphosphate 5-phosphatase; IP₃, inositol 1,4,5-trisphosphate; IP₃-3K, IP₃ 3-kinase; IP₃R, IP₃ receptor; LTD (LTP), long-term depression (potentiation); NMDA, N-methyl-D-aspartic acid; NMDAR, NMDA glutamate receptor; PLC δ (PLC β), phospholipase C δ (C β); PPR, paired-pulse ratio; PR, purinergic receptor; PSC, postsynaptic current; SIC, slow inward current; SNARE, soluble N-ethylmaleimide-sensitive factor attachment protein receptor; STDP, spike-timing-dependent plasticity; TACE, TNF α -converting enzyme; TNF α , tumor necrosis factor- α .


in the thalamus and in the cortex (Volterra and Meldolesi, 2005; Haydon and Carmignoto, 2006; Santello and Volterra, 2009; Halassa and Haydon, 2010; Navarrete et al., 2012a,b). **Figure 1A** summarizes a number of observations made at the level of hippocampal glutamatergic synapses (Bains and Oliet, 2007; Santello and Volterra, 2009). There, synaptic glutamate can trigger Ca^{2+} signaling in the surrounding astrocytic processes via metabotropic glutamate receptors (mGluRs) (Pasti et al., 1997; Fiacco and McCarthy, 2004; Panatier et al., 2011). There is also evidence in the dentate gyrus that ATP, possibly synaptically-released, triggers astrocytic Ca^{2+} signaling through the activation of metabotropic purinergic P_2Y_1 receptors (Jourdain et al., 2007; Di Castro et al., 2011; Larsson et al., 2011; Santello et al., 2011). Following elevation of intracellular Ca^{2+} , astrocytes can release glutamate as well as other chemical transmitters such as D-serine (D-ser) and ATP which can be converted into adenosine (Adn) in the extracellular milieu (Bezzi et al., 2004; Pascual et al., 2005; Montana et al., 2006; Henneberger et al., 2010; Parpura and Zorec, 2010). Astrocyte-released glutamate diffuses in the extrasynaptic space and may bind to glutamate receptors (GluRs), including mGluRs and NMDARs on neighboring presynaptic terminals, modulating the release of neurotransmitter (Fiacco and McCarthy, 2004; Jourdain et al., 2007; Perea and Araque, 2007; Bonansco et al., 2011; Di Castro et al., 2011). An analogous action on synaptic release could also be due to astrocyte-derived ATP and its derivative adenosine through presynaptic purinergic receptors (PRs), including both A_1 and A_2 receptors (Pascual et al., 2005; Halassa and Haydon, 2010; Panatier et al., 2011). On the postsynaptic side, astrocytic glutamate and D-serine may bind to extrasynaptic NR_2B -containing and postsynaptic NMDARs respectively, modulating neuronal firing and participating in the induction of long-term potentiation (Fellin et al., 2004; Bains and Oliet, 2007; Henneberger et al., 2010). Astrocyte could also release tumor necrosis factor- α (TNF α) by Ca^{2+} -dependent activation of TNF α -converting enzyme (TACE) (Bezzi et al., 2001; Santello and Volterra, 2012), which could strengthen excitatory synaptic transmission by promoting surface insertion of AMPA receptors (AMPA) (Beattie et al., 2002; Stellwagen and Malenka, 2006; Bains and Oliet, 2007). This signaling route could also play a role in pathological states such as post-traumatic epilepsy (Balosso et al., 2009; Volman et al., 2011) or spinal cord injury (Stellwagen et al., 2005; Ferguson et al., 2008). On the other hand, extracellular levels of TNF α control glutamate release from astrocytes, ultimately modulating the astrocytic action on presynaptic function (Domercq et al., 2006; Santello et al., 2011). This intricate signaling network is further complicated by the possibility that astrocyte Ca^{2+} events are triggered by additional mechanisms, including the action of ATP released extracellularly by astrocytes themselves or of IP_3 that diffuses intracellularly, from one astrocyte to another, through gap junction (GJCs) (Kang et al., 2005; Scemes and Giaume, 2006). At Schaffer collateral synapses, astrocytic Ca^{2+} increases could also be promoted by retrograde endocannabinoid signaling from postsynaptic terminals via activation of endocannabinoid CB_1 receptors (Navarrete and Araque, 2008, 2010) (omitted from **Figure 1A** for clarity). Moreover, the relation between astrocytic Ca^{2+} and gliotransmitter release is not simple: some of the Ca^{2+}

signals that can be generated in astrocytes are apparently not able to induce gliotransmitter release or its synaptic consequences (Fiacco et al., 2007; Agulhon et al., 2008, 2010; Petravicz et al., 2008; Lovatt et al., 2012), while gliotransmitters can also be released by mechanisms that are independent of Ca^{2+} signaling (Parpura and Zorec, 2010) (not included in **Figure 1A** for simplicity).

Despite its apparent complexity, the ensemble of astrocyte-synapse signaling interactions discussed above can be well captured by the modeling scheme of **Figure 1B**. This scheme shows the three essential components of astrocyte-regulated synapses, also referred to as “tripartite synapses” (Araque et al., 1999; Haydon, 2001): these are the pre- (PRE) and postsynaptic (POST) terminals, and the astrocyte, i.e., an astrocytic process surrounding the synaptic elements (ASTRO) (Araque et al., 1999; Haydon, 2001). Moreover, in addition to the classical neuronal path that leads from input presynaptic action potentials, commonly referred to as input spikes (IN), to the output postsynaptic current (OUT), further input and/or output pathways (I/O) could coexist due to the above-mentioned routes based on astrocytic Ca^{2+} signaling (Giaume et al., 2010).

Focusing on synapse-astrocyte coupling, three fundamental pathways are identified: one (**A**) from the synapse to the astrocyte whereby synaptically-released glutamate (or other synaptic agents) promotes Ca^{2+} signaling in the astrocyte and the other two (**B** and **C**) from the astrocyte to synaptic terminals, whereby glutamate or ATP released from the astrocyte affects synaptic function (Volterra and Meldolesi, 2005; Santello and Volterra, 2009). Additional pathways supported by other neuroactive agents such as D-serine or TNF α can also be evoked in parallel to those shown in **Figure 1B** but they would not alter the essence of the scheme. Moreover, although based on experimental results at excitatory synapses in the hippocampus, (Araque et al., 1998a,b; Fiacco and McCarthy, 2004; Jourdain et al., 2007; Perea and Araque, 2007; Andersson and Hanse, 2010; Santello et al., 2011), the modeling scheme in **Figure 1B** could also hold for other reported pathways such as GABA-evoked gliotransmission at interneuron-to-pyramidal cell synapses in the hippocampus (Kang et al., 1998; Serrano et al., 2006), or gliamediated ATP release at hippocampal synapses (Pascual et al., 2005), in the hypothalamus (Gordon et al., 2009), and in the retina (Newman, 2003, 2005), or glial modulation of neuromuscular transmission (Robitaille, 1998; Rousse et al., 2010; Todd et al., 2010) (see **Table 1** for a summary of the possible signaling pathways).

Analysis of the scheme in **Figure 1B** reveals that astrocytes mediate two loops in the signal flow from presynaptic to postsynaptic terminal: a *feedforward* and a *feedback* one. The feedforward loop ends on the postsynaptic terminal and is activated when synaptic glutamate and/or ATP induces glutamate and/or D-serine release from the astrocyte to the postsynaptic element, i.e. the **A–C** path in **Figure 1B** (Bains and Oliet, 2007; Barres, 2008; Santello and Volterra, 2009). The feedback loop ends on the presynaptic terminal (the **A–B** path in **Figure 1B**) and is activated when synaptic glutamate or ATP trigger Ca^{2+} -dependent release of glutamate and/or ATP from the astrocyte to the presynaptic terminal, leading to modulation of synaptic release through specific



presynaptic receptors (Santello and Volterra, 2009; Halassa and Haydon, 2010).

In principle the two pathways could coexist at the same synapse where they are expected to display different dynamics and

respond to different preferred input stimuli. Therefore their coexistence at the same synapse could give rise to complex effects that are hard to quantify when considered altogether. Accordingly, a common approach in experiments is to characterize their effects

Table 1 | Transmitters, targeted receptors, and major effects on synaptic transmission by the signaling pathways A, B, C in Figure 1B (*in situ* and *in vivo* studies only).

References	Sp. ¹	Prep. ²	Area ³	Syn. neurotr. ⁴	Targeted receptor ⁵	Gliotr. ⁶	Targeted receptors ⁵		Cell ⁷	Effects ⁸
					Pathway A		Pathway B	Pathway C		
Wang et al., 2006	M	VV	BC	Glu	mGluR-I					
Porter and McCarthy, 1996	R	ST	CA1	Glu	mGluR, iGluR					
Perea and Araque, 2007	R	ST	CA1			Glu	mGluR-I		PY	↑synaptic release probability
Liu et al., 2004a,b	R	ST	CA1			Glu	KAR		IN	↓mIPSCs (frequency)
Liu et al., 2004a,b	R	ST	HIP			Glu	mGluR-II/III		IN	↑synaptic release probability
Bardoni et al., 2010	R	ST	DH			Glu		NMDAR	SGN	SICs; synchronous Ca ²⁺ elevations in neighboring neurons
Kang et al., 2005	R	ST	CA1			Glu		iGluR	PY	SICs
Parri et al., 2001	R	ST	VBT			Glu		NMDAR	TCN	SICs
Bezzi et al., 1998	R	ST, VT	CA1, VC, COR			Glu		GluR	PY	Ca ²⁺ elevations in neighboring neurons
Newman and Zahs, 1997	R	ST	Ret ^m			Glu*		iGluR	AC*	↑neuronal inhibition
Bonansco et al., 2011	R	ST	CA1			Glu	mGluR-I	NMDAR	PY	↑mEPSCs (frequency); SICs; control of t-LTP induction
Fiacco and McCarthy, 2004	M	ST	CA1	n.r.	n.r.	Glu	mGluR-I		PY	↑synaptic release probability
Pasti et al., 1997	R	ST	CA1, VC	Glu	mGluR	Glu*			PY	Ca ²⁺ elevations in neighboring neurons
Pirttimaki et al., 2011	R	ST	VBT	n.r.	mGluR-I	Glu		NR ₂ B-NMDAR	TCN	Long-term enhancement of SIC frequency
D'Ascenzo et al., 2007	M	ST	NAcc	Glu	mGluR ₅	Glu		NR ₂ B-NMDAR	MSN	SICs, ↑neuronal firing
Fellin et al., 2004	R	ST	CA1	Glu	n.r.	Glu		NR ₂ B-NMDAR	PY	SICs; synchronous Ca ²⁺ elevations in neighboring neurons
Newman, 2005	R	ST	Ret ^m	ATP	n.r.					
Piet and Jahr, 2007	R	ST	CER ^b	Glu, ATP	AMPA, P ₂ YR					
Rieger et al., 2007	M	ST	OB	Glu, ATP	mGluR ₁ , P ₂ Y ₁ R					
Beierlein and Regehr, 2006	R	ST	CER ^b	Glu, ATP	mGluR ₁ , P ₂ YR					
Newman, 2003	R	ST	Ret ^m			ATP		A ₁ R	RN	↑K ⁺ conductance; ↑neuronal inhibition

(Continued)

Table 1 | Continued

References	Sp. ¹	Prep. ²	Area ³	Syn. neurotr. ⁴	Targeted receptor ⁵		Gliotr. ⁶	Targeted receptors ⁵		Cell ⁷	Effects ⁸
					Pathway A			Pathway B	Pathway C		
Torres et al., 2012	M	ST	CA1				ATP		P ₂ Y ₁ R	IN	↑neuronal firing
Schmitt et al., 2012	M	ST, VV	HIP	n.r.	n.r.		ATP	A ₁ R		PY	↓fEPSP (slope)
Di Castro et al., 2011	R	ST	MLDG	ATP	P ₂ Y ₁ R	n.r.	n.r.	n.r.		GC	↑synaptic release probability
Halassa et al., 2009	M	VV	COR	n.r.	n.r.		ATP	A ₁ R		CN	↓fEPSP (slope)
Jourdain et al., 2007	R	ST	MLDG	ATP*	P ₂ Y ₁ R		Glu	NR ₂ B-NMDARs		GC	↑synaptic release probability
Panatier et al., 2011	R	ST	CA1	Glu	mGluR ₅		ATP	A _{2A} R		PY	↑synaptic release probability
Schipke et al., 2008	M	ST	BC	Glu	mGluR-I		ATP	A ₁ R, P ₂ YR		PY	↑neuronal inhibition; confinement of astrocytic Ca ²⁺ signals
Pascual et al., 2005	M	VT, ST	CA1	Glu	n.r.		ATP	A ₁ R		PY	↓fEPSP (slope); heterosynaptic depression; control of long-term plasticity (metaplasticity)
Zhang et al., 2003, 2004a	R	ST, VT	CA1	Glu	n.r.		ATP	A ₁ R, P ₂ YR		PY	↓EPSP (amplitude); heterosynaptic depression
Todd et al., 2010	F	VT	NMJ	n.r.	n.r.		ATP	A ₁ R, A _{2A} R		MF	PTD (A ₁ R); PTP (A _{2A} R)
Robitaille, 1998	F	VT	NMJ				n.r.	n.r.		MF	PTP; PTD
Bowser and Khakh, 2004	M	ST	CA1	ATP, Glu	P ₂ Y ₁ R, mGluR-I		ATP		P ₂ Y ₁ R	IN	SICs, ↑neuronal firing
Gordon et al., 2009	R	ST	PVN	Glu	mGluR-I		ATP		P ₂ XR	MNC	↑mEPSCs (amplitude)
Araque et al., 2002	R	ST	CA1	ACh	mAChR						
Bélaïr et al., 2010	F	VV	NMJ	ACh, ATP	mAChR, P ₂ YR, P ₂ XR						
Navarrete et al., 2012a,b	R	ST, VV	CA1	ACh	mAChR		Glu	mGluR		PY	↑synaptic release probability; LTP
Chen et al., 2012	M	VV, ST	V1	ACh	mAChR		n.r.		NMDAR	V1N	SICs, ↑neuronal firing
Perea and Araque, 2005	R	ST	CA1	ACh, Glu	mAChR, mGluR		Glu		NMDAR	PY	SICs
Bekar et al., 2008	M	VV	COR	NE	αAR						
Kulik et al., 1999	M	ST	CER ^b	NE	α ₁ AR						

(Continued)

Table 1 | Continued

References	Sp. ¹	Prep. ²	Area ³	Syn. neurotr. ⁴	Targeted receptor ⁵		Gliotr. ⁶	Targeted receptors ⁵		Cell ⁷	Effects ⁸
					Pathway A			Pathway B	Pathway C		
Gordon et al., 2005	R	VT, ST	PVN	NE	α_1 AR		ATP		P ₂ X ₇ R	MNC	↑mEPSCs (amplitude)
Min and Nevian, 2012	R	ST	BC	ECB ^r	CB ₁ R		Glu	NMDAR		PY	↓EPSP (slope), t-LTD
Navarrete and Araque, 2010	M	ST	CA1	ECB ^r	CB ₁ R		Glu	mGluR		PY	↑synaptic release probability
Navarrete et al., 2012a,b	H	ST	HIP, COR	n.r.	GluR, PR, CBR		Glu		NMDAR	PY	SICs
Navarrete and Araque, 2008	M	ST	CA1	ECB ^r	CB ₁ R		Glu		NMDAR	PY	SICs
Panatier et al., 2006	R	ST	SON				D-ser		NMDAR	MNC	Control of long-term plasticity (metaplasticity)
Takata et al., 2011	M	VV	BC	ACh	mAChR		D-ser		NMDAR	PY	Control of LTP induction
Henneberger et al., 2010	R	ST	CA1	n.r.	n.r.		D-ser		NMDAR	PY	Control of LTP induction
Fellin et al., 2009	M	VV, ST	COR, HIP	n.r.	n.r.		D-ser		NMDAR	CN, PY	↑NMDAR-mediated currents
Yang et al., 2003	R	ST, VT	CA1	Glu	n.r.		D-ser		NMDAR	PY	Control of LTP induction
Lee et al., 2011	H	VT	COR				GABA				
Le Meur et al., 2012	R	ST	HIP				GABA, Glu		GABA _A R, NMDAR	PY	SOCs (GABA _A R); SICs (NMDAR)
Lee et al., 2010	M	ST	CER ^b				GABA		GABA _A R ^s	GC, PF ^s	↑neuronal inhibition ^s
Kozlov et al., 2006	R	ST	OB				GABA, Glu		GABA _A R, NMDAR	MC	SOCs (GABA _A R); SICs (NMDAR)
Serrano et al., 2006	R	ST	CA1	GABA	GABA _B R		ATP	A ₁ R		PY	↓fEPSP (amplitude); heterosynaptic depression
Kang et al., 1998	R	ST	CA1	GABA	GABA _B R		n.r.	iGluR	iGluR	PY	↑mIPSCs (frequency and amplitude)
Stellwagen and Malenka, 2006	M	VT	HIP				TNF α		n.r.	PY	↑AMPA; LTP
Beattie et al., 2002	R	ST, VT	HIP				TNF α		n.r.	PY	↑AMPA
Santello et al., 2011	M	ST, VT	MLDG				TNF α , Glu				Control of Glu exocytosis from astrocytes
Bezzi et al., 2001	R, H	ST, VT	HIP				TNF α , Glu				Control of Glu exocytosis from astrocytes

(Continued)

Table 1 | Continued

Reference to *in vitro* studies is included whenever the letter are missing.

1 Specimen. F, frog; H, human; M, mouse; R, rat.

2 Preparation. *VT*, *in vitro*/cultures; *ST*, *in situ*/slices; *VV*, *in vivo*.

3 Brain area or body area. *BC*, barrel cortex; *CA1*, Cornu Ammonis area 1; *CER*, cerebellum; *COR*, cortex; *DH*, dorsal horn; *HIP*, hippocampus; *MLDG*, molecular layer dentate gyrus; *NAcc*, nucleus accumbens; *NMJ*, neuromuscular junction; *PVN*, paraventricular nucleus of the hypothalamus; *Ret*, retina; *VBT*, ventrobasal thalamus; *V1*, primary visual cortex; *VC*, visual cortex.

4 Synaptically-released neurotransmitter. *Ach*, acetylcholine; *ECB*, endocannabinoids; *NE*, norepinephrine.

5 Astrocytic receptors targeted by synaptically-released neurotransmitters (pathway A) and neuronal receptors targeted by gliotransmitters (pathways B and C). α AR (α_1 AR), α - (α_1 -) adrenergic receptors; *CBR*, cannabinoid receptors; *iGluR*, ionotropic glutamate receptors; *KAR*, kainate receptor; *mACHR*, muscarinic receptors; *mGluR-I/II/III*, group I (II/III) metabotropic glutamate receptors.

6 Gliotransmitter released from astrocyte.

7 Cell targeted by gliotransmitters, where targeted receptors and effects were reported. *AC*, amacrine cell; *CN*, cortical neuron; *GC*, granule cell; *IN*, interneuron; *MF*, muscle fiber; *PF*, parallel fiber axon; *PI*, pyramidal neuron; *MNC*, magnocellular neurosecretory cell; *MSN*, medium spiny neurons; *SGN*, substantia gelatinosa neuron; *RN*, retinal neuron; *V1N*, V1 excitatory neuron

8 Effects triggered by gliotransmission on pathways B or C in **Figure 1B**. (f)EPSP, (field) excitatory postsynaptic potential; mEPSC (mIPSC), miniature excitatory (inhibitory) postsynaptic current; tLTP (t-LTD), spike-timing-dependent long-term potentiation (depression); LTP, long-term potentiation; LTD, post-tetanic depression; PTD, post-tetanic potentiation; SICS, slow inward (depolarizing) currents; SOCs, slow outward (hyperpolarizing) currents. Arrows denote increase (\uparrow) or decrease (\downarrow) and are followed by the associated signal, e.g. \uparrow (\downarrow)EPSP (frequency/amplitude/slope): increase (decrease) in (frequency/amplitude) (slope) of EPSPs; fEPSP, field EPSP.

Other: *b*, study on Bergmann glia cells; *r*, retrograde signaling; *s*, suggested by investigators; *m*, study on Müller glia cells; *n.r.*, when existence of a specific signaling pathway is observed but details are not reported; * indirect evidence.

on synaptic function by separate manipulations of pathways A–C (**Figure 1B**) using different techniques (Montana et al., 2004; Jourdain et al., 2007; Marchaland et al., 2008; Di Castro et al., 2011; Panatier et al., 2011; Santello et al., 2011). For example, characterization of the feedback pathway on the presynaptic terminal (A–B) can be carried out by opening the feedback loop (for example by inhibiting elements of A or B) and analyzing the signaling components separately. From a theoretical point of view, this approach is put forth by at least three steps of analysis that are: (1) characterization of Ca^{2+} dynamics in the astrocyte as a function of different synaptic inputs (that is, pathway A); (2) characterization of how gliotransmitter release from the astrocyte depends on different astrocytic intracellular Ca^{2+} dynamics; and (3) characterization of the effect on synaptic release of Ca^{2+} -dependent gliotransmitter release from the astrocyte (i.e. pathway B). These three aspects are discussed below from a modeling perspective, focusing on their possible roles in synaptic information processing.

CHARACTERISTICS OF ASTROCYTE Ca^{2+} EXCITABILITY AND ITS RELATIONSHIP WITH SYNAPTIC ACTIVITY

Intracellular Ca^{2+} elevations in the astrocyte are not simple on-off signals (Carmignoto, 2000; Zonta and Carmignoto, 2002; Di Castro et al., 2011). There are multiple and varied spatiotemporal patterns of Ca^{2+} elevation, which probably underlie different types of function, including generation of diverse output signals (Carmignoto, 2000; Volterra and Meldolesi, 2005; Zorec et al., 2012). Two main types of neuronal activity-dependent Ca^{2+} responses are observed in astrocytes (Grosche et al., 1999; Codazzi et al., 2001; Matyash et al., 2001; Zonta and Carmignoto, 2002; Scemes and Giaume, 2006): (1) transient Ca^{2+} increases that are confined to their distal processes (Pasti et al., 1997; Nett et al., 2002; Di Castro et al., 2011) and (2) Ca^{2+} elevations propagating along these processes as regenerative Ca^{2+} waves, often eventually reaching the cell soma (Pasti et al., 1997; Sul et al., 2004). This latter kind of response can even propagate to neighboring astrocytes, giving rise to intercellular Ca^{2+} waves (Tian et al., 2005; Kuga et al., 2011). On the other hand, intercellular Ca^{2+} propagation does not necessarily need propagation through the cell soma and has been observed across astrocytic processes or from an end foot to an end foot (Mulligan and MacVicar, 2004; Giaume et al., 2010).

The precise signaling cascades underlying the various forms of Ca^{2+} elevation are not completely understood. In general, Ca^{2+} signals in astrocytes are determined by an intricate interplay of amplification, buffering, and extrusion pathways linked to cytosolic Ca^{2+} elevations mediated by influx from the extracellular space (Malarkey et al., 2008; Shigetomi et al., 2011) and/or release from intracellular endoplasmic reticulum (ER) stores (Verkhatsky et al., 2012). IP_3 -triggered Ca^{2+} -induced Ca^{2+} release (CICR) from the ER is considered the primary mechanism responsible for intracellular Ca^{2+} dynamics in astrocytes (Volterra and Meldolesi, 2005; Nimmerjahn, 2009). This mechanism, schematized in **Figure 2A**, is essentially controlled by the interplay of three fluxes: (1) a Ca^{2+} transfer from the cytosol to the ER (J_P) mediated by endoplasmic-reticulum Ca^{2+} -ATPase (SERCA) pumps which contributes to the maintenance of higher

Ca^{2+} concentrations in the ER stores than in the cytosol; (2) a passive Ca^{2+} leak (J_L) from the ER to the cytosol that is driven by the Ca^{2+} gradient between the ER and the cytosol; and (3) an efflux (J_{NL}) from the ER to the cytosol through IP_3 receptor (IP_3R) channels, which depends both on IP_3 and Ca^{2+} concentrations in the cytosol in a nonlinear fashion (Bezprozvanny et al., 1991; Ramos-Franco et al., 2000; Shinohara et al., 2011).

Cytosolic Ca^{2+} regulates IP_3Rs in a biphasic manner: Ca^{2+} release from the ER is potentiated at low cytosolic Ca^{2+} concentrations but is inhibited at higher Ca^{2+} concentrations (Iino, 1990; Bezprozvanny et al., 1991). On the other hand, IP_3 monotonically activates IP_3R channels at constant Ca^{2+} concentrations (Watras et al., 1991), but dynamically changes the Ca^{2+} sensitivity of the channel (Kaftan et al., 1997; Ramos-Franco et al., 2000; Mak et al., 2003). At low, subsaturating IP_3 concentrations, the optimal Ca^{2+} concentration for IP_3R modulation becomes lower, whereas at very high IP_3 concentrations, channel activity persists at supramicromolar Ca^{2+} concentrations (Kaftan et al., 1997; Mak et al., 2003). Thus, the level of IP_3 determines the dynamics of intracellular Ca^{2+} .

Both production and degradation of IP_3 depend on enzymes that are regulated by cytosolic Ca^{2+} (Berridge et al., 2003; De Pittà et al., 2009a,b). These include Ca^{2+} -dependent PLC δ -mediated IP_3 synthesis and Ca^{2+} -dependent IP_3 degradation by IP_3 3-kinase (IP_3 -3K) and by inositol polyphosphate 5-phosphatase (IP -5P) (Figure 2A) (Zhang et al., 1993; Sims and Allbritton, 1998; Rebecchi and Pentylala, 2000). However, while the activity of IP_3 -3K is stimulated by cytosolic Ca^{2+} (Communi et al., 1997), IP -5P is inhibited instead (Communi et al., 2001). This results in different mechanisms of IP_3 degradation depending on the Ca^{2+} concentration in the cytoplasm (Sims and Allbritton, 1998; Irvine et al., 2006). Thus, for example, for equally-expressed enzymes, at low Ca^{2+} concentrations, namely lower than 500 nM (Sims and Allbritton, 1998; De Pittà et al., 2009a,b), IP_3 degradation is promoted by both IP -5P and IP_3 -3K, whereas for intermediate-to-high cytosolic Ca^{2+} concentrations, degradation by IP_3 -3K becomes predominant (Sims and Allbritton, 1998). Theoretical investigation showed that the interplay of these two regimes is both necessary and sufficient to reproduce Ca^{2+} oscillations and pulsations observed experimentally (De Pittà et al., 2009a,b).

Intracellular levels of IP_3 can also be controlled by gap junction mediated diffusion of IP_3 from other regions of the same astrocyte or from neighboring cells (Giaume et al., 2010) (i.e., the I/O pathway in Figure 1B). Moreover, synaptic glutamate (or other synaptic agents) can bind to astrocytic G protein-coupled receptors (GPCRs) like mGluRs that are directly linked to intracellular IP_3 production by PLC β (Zur Nieden and Deitmer, 2006) (pathway A in Figure 1B). In this fashion, astrocytic Ca^{2+} dynamics triggered by synaptically-controlled IP_3 production can be regarded as a form of encoding information about activity in neighboring synapses.

Encoding of synaptic activity by astrocytic Ca^{2+} is likely multimodal, depending on many possible intracellular properties (De Pittà et al., 2008, 2009a,b; Dupont et al., 2011). A widely adopted classification considers the amplitude and the frequency of Ca^{2+} increases from resting levels (Berridge, 1997; Falcke, 2004; De Pittà et al., 2008, 2009a,b). In this view, as summarized

in Figure 2B, synaptic activity reflected by different intracellular IP_3 concentrations (STIMULUS), is encoded by the modulation of Ca^{2+} oscillations and pulsations either in their amplitude (AM), their frequency (FM), or both (AFM). While available experimental data suggest a preferential FM mode of encoding (Pasti et al., 1997), AM and AFM encoding of synaptic activity are also plausible mechanisms given that the amplitude of Ca^{2+} response can strongly depend on the stimulation intensity (Wang et al., 2006; Di Castro et al., 2011; Panatier et al., 2011; Torres et al., 2012). This is the case for example, of synaptic inputs that occur rapidly one after the other whereby the ensuing intracellular Ca^{2+} concentration builds up as a cumulation of such inputs (Perea and Araque, 2005; Torres et al., 2012). In further support of the AM/AFM encoding is the experimental observation that glutamate exocytosis from the astrocyte occurs only when Ca^{2+} increases beyond a threshold concentration (Newman and Zahs, 1997; Parpura and Haydon, 2000; Pasti et al., 2001; Auld and Robitaille, 2003; Montana et al., 2006). Hence, astrocytic Ca^{2+} increases in response to synaptic activity would not systematically trigger the release of glutamate or other gliotransmitters from the astrocyte. Acting on the amplitude of astrocytic Ca^{2+} signals, AM/AFM encodings could constitute a way to regulate astrocytic gliotransmitter release by synaptic activity. Further experiments are needed to elucidate the nature of the Ca^{2+} threshold for astrocytic exocytosis since this latter might be gliotransmitter-specific (Montana et al., 2006). Accordingly, AM/AFM encoding of Ca^{2+} dynamics could vary from one gliotransmitter to another.

Experimental evidence suggests that Ca^{2+} dynamics does not simply mirror synaptic activity but is more complex, to a point that astrocytes have been proposed to perform genuine processing of synaptic information (Perea and Araque, 2005; Perea et al., 2009). This possibility follows from the complex network of IP_3 and Ca^{2+} signaling and subtends a scenario where Ca^{2+} could be only one of the players in the encoding and processing of synaptic activity by astrocytes (Mishra and Bhalla, 2002). Yet many, if not all, of the other signals underlying the complex cascade of biochemical reactions that link synaptically-released glutamate to CICR, could also carry out encoding and processing (Barlow, 1996; Berridge et al., 2003). Theoretical investigations suggested that IP_3 could also encode for the glutamate stimulation levels via a systematic AFM encoding (De Pittà et al., 2009a,b) as shown in Figure 2C. When cytosolic Ca^{2+} levels are low, close to resting values, IP_3 generally increases with ongoing synaptic activity (Glu). With low Ca^{2+} , the activity of IP_3 -3K is reduced and the resulting IP_3 degradation slows down. The contribution to IP_3 production by Ca^{2+} -dependent PLC δ is reduced as well, so that intracellular IP_3 mostly depends on the frequency of synaptic release. Rapid successions of synaptic release events produce crisp increases of IP_3 (essentially proportional to the number of successive synaptic release events) while, between two remote release events, IP_3 tends to relax to resting levels. As a result, IP_3 dynamics overall evolves as the integral of synaptic activity. If IP_3 reaches the CICR-triggering threshold, intracellular Ca^{2+} increases fast and so does IP_3 -3K activity. Then, IP_3 is rapidly degraded and resting IP_3 levels are restored, thus resetting the integral of synaptic activity to initial values.




FIGURE 2 | Computational aspects of astrocytic Ca $^{2+}$ signaling.

(A) Scheme of IP $_3$ -mediated Ca $^{2+}$ -induced Ca $^{2+}$ release in the astrocyte. Calcium and IP $_3$ signals are controlled by synaptic glutamate through metabotropic glutamate receptor- (mGluR-) PLC β -mediated IP $_3$ production (see text for details). (B) IP $_3$ and Ca $^{2+}$ signals can be envisioned to encode incoming synaptic activity through frequency and amplitude of their oscillations. Astrocytes could thus encode synaptic information either by modulations of the amplitude (AM), the frequency (FM), or both (AFM) of their Ca $^{2+}$ oscillations. (C) Simulated Ca $^{2+}$ and IP $_3$ patterns in response to sample synaptic glutamate release (Glu) in a model astrocyte (De Pittà et al., 2009a,b) reveals that IP $_3$ signals could be locked in the AFM-encoding independently of the encoding mode of the associated Ca $^{2+}$ signals. This feature could allow the astrocyte to optimally integrate synaptic stimuli.

(D, top) Simulated Ca $^{2+}$ (black trace) and IP $_3$ (green trace) signals in the same astrocyte model as in (C), and associated rates of IP $_3$ production (prod.) and degradation (degr.) (middle panels, dashed black lines) in response to two consecutive synaptic glutamate release events (bottom panel, Glu). The analysis of the contributions of different enzymes to IP $_3$ signaling (solid colored traces; cyan: PLC β ; pink: PLC δ ; orange: IP-5P; and purple: IP $_3$ -3K) reveals dynamical regulation by Ca $^{2+}$ of different mechanisms of IP $_3$ production/degradation which could ultimately underlie dynamical regulation of astrocyte processing of synaptic stimuli. (E) Simulated propagation of Ca $^{2+}$ waves in a heterogeneous linear chain composed of both FM (red traces) and AFM (green traces) astrocytes reveals that encoding of synaptic activity (STIMULUS) could change according to cell location along the chain. Adapted from Goldberg et al. (2010).

It is precisely the alternation between these two different phases of IP $_3$ degradation (a high Ca $^{2+}$ —high IP $_3$ -3K-activity phase and a low Ca $^{2+}$ —low IP $_3$ -3K-activity phase), that endows IP $_3$ signal with high amplitude variability. On the other hand, such AM features still allow fast variations, thus rich spectral content (i.e., FM features), in response to changes in frequency of

synaptic release. This enticing possibility could endow the IP $_3$ signal with the necessary properties to function as optimal interface between synaptic stimuli and intracellular Ca $^{2+}$ signals. Since neural information is carried by the timing of spikes rather than by their amplitude (Sejnowski and Paulsen, 2006), the capability of fast highly-variable amplitude changes corresponding to

rich spectral content of IP₃ signals, would fulfill this requirement, embedding the essential spectral features of the synaptic signal into the spectrum of the IP₃ transduction. On the other hand, because Ca²⁺ signals are triggered primarily by suprathreshold IP₃ elevations (Li et al., 1994; Keizer et al., 1995), the coexistence of AM features within the AFM IP₃ signal seems to be a necessary prerequisite in order to trigger CICR.

This could also help elucidate the origin of the integrative properties of Ca²⁺ signaling in astrocytes (Perea et al., 2009). These properties could result from at least two steps of integration: one step is the transduction of the agonist signal into the IP₃ signal; the other step is the cross-talk between IP₃ and Ca²⁺ signals. Hence, AFM-encoding IP₃ dynamics could operate a first preliminary integration by smoothing the highly indented synaptic stimulus. The inherent features of CICR would then bring forth a further integration step, yielding Ca²⁺ patterns that are even smoother than IP₃ signals (De Pittà et al., 2009a,b).

The tight dynamical coupling between IP₃ and Ca²⁺ signals also suggests that the way astrocytes process synaptic signals, i.e. their frequency response to synaptic stimuli, is not fixed but rather dynamical and dependent on the history of activation of the astrocyte. This is because different IP₃ signaling mechanisms that are dynamically regulated by Ca²⁺ likely correspond to different frequency responses of the astrocyte with respect to synaptic signals. **Figure 2D** shows the time course of IP₃ production (IP₃ prod.) and degradation (IP₃ degr.) (*dashed black lines*) underlying simulated IP₃ and Ca²⁺ signals (*top panel, green and black traces*, respectively) in response to two events of synaptic glutamate release (*bottom panel*). While Ca²⁺-dependent IP₃ production by PLCβ (*cyan trace*) and PLCδ (*pink trace*) could modulate the threshold frequency of synaptic stimuli that triggers Ca²⁺ signaling in the astrocyte, existence of different regimes of IP₃ degradation within a single Ca²⁺ oscillation cycle could be responsible for different cutoff frequencies of synaptic release beyond which Ca²⁺-mediated astrocyte processing of synaptic stimuli ceases. In particular, the cutoff frequency during low IP₃-3K activity could be mainly set by the rate of IP₃ degradation by IP-5P (*orange trace*). When Ca²⁺ is high instead, IP₃ degradation by IP₃-3K (*purple trace*) could also become very strong, thus sensibly reducing the cutoff frequency. That is, the cutoff frequency is dependent on the context of underlying Ca²⁺ signaling which, in turn, depends on the history of activation of the astrocyte by synaptic stimuli.

The subcellular arrangement of the enzymes underlying IP₃ signaling could also be responsible for spatial heterogeneity of the frequency response of the astrocyte. Although the subcellular localization of IP₃ production and degradation enzymes in astrocytes remains to be elucidated, studies in brain tissue suggest that PLCβ and IP-5P could localize mainly in proximity to the plasma membrane, whereas PLCδ and IP₃-3K are preferentially in the cytoplasm (Rebecchi and Pentylala, 2000; Irvine et al., 2006). Given that the ER distribution changes from astrocytic processes to soma (Pivneva et al., 2008), different subcellular regions of the astrocyte could correspond to different cytoplasmic volumes and thus to distinctly different expressions of enzymes mediating IP₃ signaling. The ensuing different subcellular arrangement of these enzymes could ultimately provide

anatomical specificity to the astrocytic phosphoinositide signaling which underlies CICR-based astrocyte processing of synaptic activity (Fukaya et al., 2008). In this fashion, processing of synaptic stimuli by Ca²⁺ signaling at astrocytic processes could differ from that carried out in the soma by means of differently expressed IP₃ signaling-related enzymes.

COMPUTATIONAL ASPECTS OF PROPAGATING Ca²⁺ SIGNALS

Intracellular and intercellular propagation of Ca²⁺ could contribute new encoding and processing modes, in addition to those depicted in **Figure 2B**. However, despite the numerous modeling studies developed to account for the rich dynamics of astrocyte Ca²⁺ signaling (Bennett et al., 2008; Goldberg et al., 2010; Dupont et al., 2011); [for a recent review on calcium modeling see Falcke (2004)], we still lack a comprehensive theoretical framework to link the local Ca²⁺ signals that are restricted to small regions of the astrocytic processes, to their spatial dynamics and their possible propagation at larger spatial scales: intracellular propagation, global whole-cell signals or cell-to-cell Ca²⁺ waves.

Propagations at these various scales probably differ by their underlying mechanisms (Falcke, 2004; Scemes and Giaume, 2006). Fast-rising and short-lived local Ca²⁺ events, observed in response to even a single quantal release from synaptic terminals (Di Castro et al., 2011; Panatier et al., 2011) closely resemble spatially confined Ca²⁺ puffs or blips in other cell types (Thomas et al., 2000; Bootman et al., 2001) and could depend on spatial clustering of IP₃Rs along the ER structures or of mGluRs along the plasma membrane, or on both (Marchaland et al., 2008; Panatier et al., 2011; Arizono et al., 2012). The cumulative recruitment of these Ca²⁺ puffs could lead to spatially more extended Ca²⁺ events which could either be still confined within astrocytic processes (Di Castro et al., 2011) or propagate to other cellular regions or to other cells as regenerative Ca²⁺ waves (Pasti et al., 1997; Kuga et al., 2011).

Calcium could propagate by at least two routes [for a recent review see Scemes and Giaume (2006)]. One is intracellular, through GJCs, involving diffusion of IP₃ directly from cytoplasm to cytoplasm. The other route is extracellular, involving release of ATP from the astrocyte which binds to GPCRs of the same cell or neighboring astrocytes, increasing their IP₃ levels (Guthrie et al., 1999). The relative contribution of each of these pathways likely depends on developmental, regional and physiological states and could subtend different ranges of propagations as well as different temporal features (Haas et al., 2006; Scemes and Giaume, 2006; Giaume et al., 2010).

The restriction and clustering of mGluRs expression along astrocytic processes to subregions that colocalize with synaptic terminals (van den Pol et al., 1995; Arizono et al., 2012) hints the possibility of a subcellular compartmentalization of Ca²⁺ signals (Marchaland et al., 2008; Di Castro et al., 2011; Panatier et al., 2011). Local Ca²⁺ events would be spatially restricted to narrow regions around each mGluRs cluster thus defining independent signals within the same process. In this fashion, astrocytes could carry out parallel integration and processing of synaptic information on different temporal and spatial scales at different processes (Goldberg et al., 2010; Bernardinelli et al., 2011) or even in

different subregions of the same process which could constitute separate functional microdomains (Panatier et al., 2011). Furthermore, the resulting neuromodulatory action exerted by astrocytic gliotransmitters on synaptic terminals and their impact on neuronal network activity could vary from one process or microdomain to the other (Navarrete and Araque, 2011).

The spatial scale and the time window of Ca^{2+} events likely discriminate between different mechanisms of IP_3 and Ca^{2+} signaling. At the level of astrocytic processes and subcellular compartments, Ca^{2+} propagation could be mediated by fast intracellular IP_3 linear (i.e. Fickian) diffusion (Sneyd et al., 1994; Falcke, 2004). On the other hand, when considering intercellular propagation mediated by gap junctions, IP_3 transport from one cell to the other could be essentially nonlinear. This scenario was shown to be consistent with the observed variability of Ca^{2+} wave propagation distance (Goldberg et al., 2010) and could explain the long-range regenerative propagation of Ca^{2+} waves observed in cortical astrocytes (Scemes and Giaume, 2006; Tian et al., 2006).

Cell heterogeneity likely constitutes an additional critical aspect that substantially affects propagation patterns and extent of propagation of Ca^{2+} signals (Iacobas et al., 2006; De Pittà et al., 2008; Goldberg et al., 2010). This possibility is illustrated in **Figure 2E** by a toy example consisting of a heterogeneous linear chain of FM-encoding (*red traces*) and AFM-encoding (*green traces*) astrocytes. The synaptic stimulus is restricted to the first cell of the chain (i.e. cell number 1). Only FM-encoding cells guarantee regenerative propagation of Ca^{2+} signals, whereas AFM cells do not, acting like propagation barriers along the chain. Unlike in AFM cells, Ca^{2+} -dependent IP_3 production in FM cells guarantees IP_3 diffusion to the next cell in the chain to levels that are beyond the threshold of CICR thus promoting regenerative propagation (Goldberg et al., 2010). Moreover, the shape of the local Ca^{2+} events in each cell changes along the propagation path: after each AFM cell, the frequency of FM-encoding Ca^{2+} pulses is reduced, suggesting that different propagation patterns could carry out different processing of synaptic information (Goldberg et al., 2010).

The above scenario hints that the spatial distribution of astrocytes in different brain areas could be made to fulfill specific processing tasks. Indeed neighboring astrocytes in the brain are believed to be distributed in space in a non-random orderly fashion called “contact spacing” (Chan-Ling and Stone, 1991; Volterra and Meldolesi, 2005) or “tiling,” where each astrocyte creates its micro-anatomical domain with its processes overlapping with adjacent astrocytes only at their periphery (Bushong et al., 2002). Such spatial arrangement, combined with the heterogeneity of astrocytic responses, could be important in intercellular Ca^{2+} wave propagations and the related computational tasks carried out by astrocyte networks. The latter could be relevant in particular for the emergence of astrocytic functional maps observed in several brain areas (Giaume et al., 2010).

Increasing evidence argues in fact for a functional organization of astrocytes, reminiscent of that of cortical neurons (Bernardinelli et al., 2011). In the ferret visual cortex astrocytes, like neurons, respond to visual stimuli, with distinct spatial receptive fields and sharp tuning to visual stimulus features, including orientation and spatial frequency (Schummers et al., 2008). The

stimulus-feature preferences of astrocytes there can be mapped across the cortical surface, in close register with neuronal maps (Schummers et al., 2008). Similar observations were also reported for astrocytes in the motor cortex (Haas et al., 2006) and in the somatosensory cortex (Schipke et al., 2008) as well as in the olfactory bulb (De Saint Jan and Westbrook, 2005).

Anatomical compartmentalization of astrocyte networks could underlie such functional organization. Both in the somatosensory barrel cortex and in the olfactory bulb, astrocytes are preferentially connected by gap junctions within the same barrel or olfactory glomerulus rather than between adjacent barrels or glomeruli (Houades et al., 2008; Roux et al., 2011). However, spatial confinement of gap junctions within single barrels/glomeruli might not fully account for the selective activation of astrocyte by electrical stimulation. Indeed, similar Ca^{2+} signals in response to the same stimulus could be observed in the same barrel yet with pharmacological block of astrocyte gap junctions (Schipke et al., 2008). Thus, additional factors must contribute to the astrocyte tuned response, which likely emerges as a result of the dynamical interactions with surrounding neurons (Rouach et al., 2004) and could ultimately depend on location and nature of activated cells among the other astrocytes in the network (Matyash and Kettenmann, 2010; García-Marqués and López-Mascaraque, 2012). In the case of **Figure 2E**, for example, stimulated FM-encoding cells could trigger Ca^{2+} signaling in neighboring cells in a regenerative fashion thus extending their tuned response in space. On the other hand, AFM cells, acting as propagation barriers, could shape the borders of this tuned response, eventually drawing the topographical features of the ensuing functional map (Lallouette and Berry, 2012).

RELATING Ca^{2+} SIGNALS TO GLIOTRANSMITTER EXOCYTOSIS FROM THE ASTROCYTE

There is a number of possible routes by which astrocytes could release gliotransmitters (Ni et al., 2007; Parpura and Zorec, 2010; Zorec et al., 2012), but Ca^{2+} -dependent exocytosis is likely the major one on a physiological basis (Barres, 2008; Parpura et al., 2011). However, the identity of incoming inputs, the underlying molecular mechanism and the physiological conditions that govern gliotransmitter exocytosis largely remain to be elucidated (Montana et al., 2006; Ni et al., 2007; Santello and Volterra, 2009; Parpura and Zorec, 2010). Calcium-dependent exocytosis of glutamate or ATP from astrocytes, for example, may strictly depend on the nature of the upstream Ca^{2+} signal (Perea and Araque, 2005; Li et al., 2008; Marchaland et al., 2008; Pryazhnikov and Khiroug, 2008; Malarkey and Parpura, 2011); including the type of neurotransmitter involved and the type of receptor engaged (Enkvist and McCarthy, 1992; Muyderman et al., 2001; Coco et al., 2003; Bezzi et al., 2004; Blomstrand and Giaume, 2006). The influence of astrocytic glutamate or ATP on synaptic activity also likely depends both on the type of incoming stimulus and on the specific localization of the engaged receptor in the astrocyte (Perea and Araque, 2005; Santello and Volterra, 2009).

A large amount of evidence suggests that gliotransmitter exocytosis from astrocytes bears several similarities with its synaptic homologous (Bergersen and Gundersen, 2009; Santello and Volterra, 2009; Bergersen et al., 2012) (**Figure 3A**). Astrocytes




FIGURE 3 | Linking gliotransmitter exocytosis to various Ca²⁺ encoding modes. (A) Calcium-dependent glutamate and ATP exocytosis from astrocytes are both brought forth by a vesicular compartment in the astrocyte competent for regulated exocytosis. The frequency of exocytotic events is directly controlled by the shape and frequency of Ca²⁺ oscillations. (B) Modeling concept for an "exocytosis event" from the astrocyte. Calcium (*top trace*) triggers exocytosis of glutamate or ATP every time it increases beyond a certain

threshold concentration value (*red dashed line*). The overall release can then be approximated, under proper assumptions, by an exponentially-decaying pulse of extracellular concentration of glutamate or ATP (*bottom trace*). (C) Distinct Ca²⁺ encoding patterns could translate into distinct rates of gliotransmitter exocytosis events. In this way, synaptic activity encoded by astrocytic Ca²⁺ signals is linked to the frequency of glutamate/ATP release from the astrocyte in a unique fashion. Adapted from De Pittà et al. (2011).

possess vesicular compartments that are competent for the regulated exocytosis of glutamate (Bezzi et al., 2004; Bergersen and Gundersen, 2009) and ATP (Coco et al., 2003; Jaiswal et al., 2007; Zhang et al., 2007). Similarly to synapses, astrocytes express soluble N-ethylmaleimide-sensitive factor attachment protein receptors (SNARE) necessary for exocytosis (Parpura et al., 1995; Schubert et al., 2011) as well as proteins responsible for concentrating glutamate or ATP into vesicles (Bezzi et al., 2004; Montana et al., 2004; Zhang et al., 2004b; Sawada et al., 2008). Fusion with the plasma membrane, trafficking and recycling of astrocytic glutamate and ATP secretory vesicles have been observed (Bezzi et al., 2004; Chen et al., 2005; Crippa et al., 2006; Jaiswal et al., 2007; Pangršić et al., 2007; Stenovec et al., 2007), which are indicative of quantal glutamate and ATP release (Del Castillo and Katz, 1954; Pasti et al., 2001; Domercq et al., 2006; Jaiswal et al., 2007; Pangršić et al., 2007; Marchaland et al., 2008; Santello et al., 2011).

Glutamate exocytosis from cultured astrocytes evoked by GPCRs is observed with short delay, i.e. 50–100 ms, after the rise in submembrane Ca²⁺, and is mediated by a rapid succession of fusion events which peaks within ~500 ms from the onset and decays to baseline much more slowly (>1 s), though generally before the recovery of basal Ca²⁺ levels (Domercq et al., 2006; Marchaland et al., 2008; Santello et al., 2011) (refer also to **Table 2**). In addition to this "exocytotic burst," larger delays with slower rate of sustained vesicular fusion events have also been reported (Kreft et al., 2004; Malarkey and Parpura, 2011). The delayed onset with respect to the Ca²⁺ rise is consistent with a threshold Ca²⁺ concentration for release (Pasti et al., 1997; Parpura and Haydon, 2000). On the other hand the overall release of glutamate is characterized by a rising phase much faster than its decay and can be approximated by an exponential function like synaptic exocytosis yet with a decay time much slower than the latter (Marchaland et al., 2008; De Pittà et al., 2011; Santello et al.,

Table 2 | Comparison of time scales of rise, decay and full-width half-maximum (FWHM) duration of changes of intracellular Ca^{2+} , extracellular glutamate (Glu) and ATP in astrocytes and neurons.

Signal	Origin	τ_{rise} [s]	τ_{decay} [s]	FWHM [s]	Essential references
Ca^{2+}	Astrocyte (soma)	2–20 ^f	3–25 ^f	5–160	Hirase et al., 2004; Nimmerjahn et al., 2004; Wang et al., 2006
	Astrocyte (soma) ^a	~0.5	~1.1	~2–4	Winship et al., 2007
	Neuron (soma)	2–5·10 ⁻³	0.1–0.4	~0.1–0.3	Svoboda et al., 1997; Waters et al., 2003; Nimmerjahn et al., 2004
	Astrocyte (process)	0.1–0.2	0.2–4 ^f	~0.5–4	Di Castro et al., 2011; Panatier et al., 2011
	Neuron (presynaptic bouton)	0.5–5·10 ⁻³	~0.1–2	0.1–1*	Regehr et al., 1994; Emptage et al., 2001
Glu	Astrocyte	0.2–0.5	0.5–1.5	1–6	Domercq et al., 2006; Marchaland et al., 2008; Santello et al., 2011
	Neuron (synapse)	1–5·10 ⁻⁴	0.01–0.1	0.01–0.1*	Raghavachari and Lisman, 2004; Herman and Jahr, 2007; Okubo et al., 2010
ATP	Astrocyte	0.1–0.5	2–3	2–20	Pangršić et al., 2007; Li et al., 2008
	Neuron (synapse)	1–5·10 ⁻⁴ *	0.2–1*	0.15–0.5	Dundwiddie and Masino, 2001; Pankratov et al., 2007

Glutamate and ATP values refer to transient increases of their extracellular concentrations following release in a Ca^{2+} -dependent fashion. Therefore they describe the time course of the overall glutamate and ATP released by an exocytotic burst rather than by a single exocytotic event which can be much faster and occur within the first ~50 ms from Ca^{2+} rise (see text). Indicative rise and decay time constants as well as FWHM values are reported in terms of min–max ranges. Fast calcium signals imaging in astrocyte in the somatosensory cortex reported by Winship et al. (2007) are reported separately and dubbed by “a.” The letter “f” stands for values that were obtained by fitting of experimental data by a biexponential function such as $f(t) = C \cdot (\exp(-t/\tau_{\text{decay}}) - \exp(-t/\tau_{\text{rise}}))$ with C being a proper scaling factor. Asterisk “*” denotes values estimated by a model of astrocytic and synaptic release introduced in De Pittà et al. (2011).

2011) (**Figure 3B**). Although much less characterized than its glutamate counterpart, astrocytic ATP exocytosis could occur in a qualitatively similar fashion in spite of different underlying kinetics (Pangršić et al., 2007; Pryazhnikov and Khiroug, 2008; Li et al., 2008). Based on these arguments, it was proposed that astrocytic gliotransmitter release could be modeled using the same mathematical description of synaptic release, although the two mechanisms are likely different in their molecular machinery, with the kinetics of astrocyte release much slower than synaptic release (De Pittà et al., 2011; Schubert et al., 2011).

How are different modes of Ca^{2+} encoding translated into glutamate or ATP release? Assuming proper conditions about the rate of clearance of these gliotransmitters with respect to the underlying intracellular Ca^{2+} dynamics that mediate their exocytosis (Abbracchio et al., 2009), an intriguing theoretical prediction is that various patterns of Ca^{2+} oscillations could mostly correspond to different rates of gliotransmitter release (De Pittà et al., 2011). This is presented in **Figure 3C** where, for three stereotypical patterns of Ca^{2+} oscillations, that is AM, FM, and AFM, the corresponding timing of gliotransmitter release from the astrocyte is shown. In this example, FM-encoding Ca^{2+} oscillations always cross the threshold for exocytosis (*dashed red line*), triggering gliotransmitter release every time. Conversely, AM or AFM oscillations may not be large enough to reach such threshold, resulting in some oscillations that fail to trigger gliotransmitter release. In this fashion, while FM Ca^{2+} oscillations trigger gliotransmitter exocytosis at their own frequency, the amplitude of AM and AFM oscillations could selectively discriminate which Ca^{2+} pulse triggers exocytosis, eventually dictating the frequency of “measured” glutamate or ATP release events (De Pittà et al., 2011). Further experimental investigations are required to elucidate whether such prediction could effectively mimic reality.

While astrocytic Ca^{2+} signals could be triggered both by spontaneous and evoked synaptic release, gliotransmitter release might be not (Di Castro et al., 2011; Panatier et al., 2011). Blockade of Ca^{2+} -dependent glutamate release in astrocytes in the dentate gyrus was indeed observed to be effective in reducing the frequency of synaptic release events only when these were evoked by action potentials, but not when they happened spontaneously. This suggests that release of glutamate from astrocytes in this region could effectively occur only in presence of evoked synaptic activity (Di Castro et al., 2011). Interestingly, the Ca^{2+} elevations in astrocyte processes triggered by action potentials were reported to be more complex, larger in amplitude and more extended in space, than those generated by spontaneous synaptic release events. Since glutamate could be released from several sites along the same astrocyte process and this latter could contact several synapses (Panatier et al., 2011), one of such Ca^{2+} elevations could then generate multiple spatially-distinct glutamate release events modulating synaptic release at several other synapses. The same Ca^{2+} elevation however, could bring forth at subcellular regions of the process, different local Ca^{2+} dynamics, in close analogy to the behavior of the heterogeneous AM/AFM oscillations in the astrocyte chain in **Figure 2E**. Such different local Ca^{2+} dynamics could in turn result in different rates of glutamate release and thus in different modulations of synaptic release in a non-random fashion. In this way, the same astrocytic process or a segment of it, could carry out multiple regional modulations of synaptic release, depending both on the temporal and spatial dynamics of synaptic release. It is however possible that the mechanisms underlying Ca^{2+} -dependent gliotransmission differ in different brain areas (Matyash and Kettenmann, 2010; Zhang and Barres, 2010). Indeed, in contrast with the above observations in the dentate gyrus, astrocytes in the stratum radiatum

of the hippocampus were reported to release glutamate even in the absence of synaptic activity evoked by action potentials (Bonansco et al., 2011).

Another issue is what makes a single astrocyte release one gliotransmitter rather than another. A possibility is that different gliotransmitters are released in response to different stimuli. *In situ* studies indeed suggest that purinergic GPCR-mediated astrocytic Ca^{2+} signals could preferentially trigger glutamate release (Jourdain et al., 2007; Perea and Araque, 2007; Di Castro et al., 2011) (see also **Table 1**). In contrast, Ca^{2+} elevations triggered by glutamate could bring forth ATP release (Pascual et al., 2005; Gordon et al., 2009; Panatier et al., 2011). Overall these observations hint that a tight association likely exists between the type of targeted astrocytic receptor and the secretory machinery of gliotransmitters (Zorec et al., 2012). Moreover, in the same astrocyte, different gliotransmitters could be contained in different organelles with different secretory properties in response to Ca^{2+} signals (Coco et al., 2003). Indeed, while glutamate seems to be preferentially released by synaptic-like microvesicles (Bezzi et al., 2004; Jourdain et al., 2007; Bergersen et al., 2012), ATP is likely released by dense-core granules (Coco et al., 2003), and/or lysosomes (Jaiswal et al., 2007; Zhang et al., 2007; Li et al., 2008). Although the underlying molecular machinery of exocytosis remains to be elucidated, each organelle population is likely secreted in a different fashion (Pryazhnikov and Khiroug, 2008). *In vitro* evidence showed in fact that Ca^{2+} signals that triggered release of glutamate-containing vesicles (Marchaland et al., 2008) did not release ATP-filled organelles and vice versa (Coco et al., 2003; Li et al., 2008), ultimately suggesting that glutamate and ATP could be released in response to different Ca^{2+} signals (Parpura and Zorec, 2010).


Intriguingly, stimulation of astrocyte GPCRs can evoke fast gliotransmitter exocytosis within few hundreds of milliseconds (Bezzi et al., 2004; Domercq et al., 2006; Marchaland et al., 2008; Santello et al., 2011), indicating that the Ca^{2+} -dependent process that couples stimulus with secretion must be fast. Indeed, the peak of GPCR-mediated Ca^{2+} release from the ER can be as fast as 50–250 ms (Marchaland et al., 2008; Di Castro et al., 2011; Panatier et al., 2011) and Ca^{2+} -dependent exocytosis of single glutamate vesicles can occur within less than 5–20 ms from Ca^{2+} elevation (Chen et al., 2005; Bowser and Khakh, 2007; Marchaland et al., 2008; Santello et al., 2011). Therefore, to assure fast stimulus-secretion coupling, IP_3 diffusion from the site of production by GPCRs at the plasma membrane to IP_3 receptors on the ER membrane must also be of the order of tens of milliseconds and so must Ca^{2+} diffusion from the mouth of IP_3 Rs to the Ca^{2+} sensor of exocytosis (Zhang et al., 2003, 2004a; Ni et al., 2007). Given that the diffusion times of IP_3 or Ca^{2+} can be estimated as the half of the square of the diffusion distance from the site of their production in the cytoplasm divided by the respective diffusion constants (Syková and Nicholson, 2008), which can be as high as $\sim 200\text{--}300\ \mu\text{m}^2/\text{s}$ for IP_3 (Allbritton et al., 1992; Sneyd et al., 1994) and $30\ \mu\text{m}^2/\text{s}$ for Ca^{2+} (Kang and Othmer, 2009), then diffusion times of the order of milliseconds could be obtained only for diffusion lengths at most in the micron range. This prediction is fully supported by the functional evidence of local GPCR-evoked, ER-dependent Ca^{2+} microdomains

in astrocytic processes (Di Castro et al., 2011; Panatier et al., 2011) and by the morphological evidence that at astrocytic processes the ER stores are found at $\sim 200\text{--}600\ \text{nm}$ distance both from the plasma membrane and gliotransmitter-containing vesicles (Bezzi et al., 2004; Marchaland et al., 2008; Bergersen et al., 2012). Moreover, GPCR agonist-mediated Ca^{2+} transients were observed restricted areas beneath the plasma membrane, in close proximity to the sites of exocytosis (Marchaland et al., 2008), suggesting that both GPCRs and secretory vesicles must be spatially close to the IP_3 Rs responsible for Ca^{2+} release from the ER. This indicates that the location of IP_3 Rs along the ER membrane in astrocyte processes might be non-random (Blaustein and Golovina, 2001), in tight spatial association with astrocytic GPCRs on the plasma membrane and releasable gliotransmitter organelles (Marchaland et al., 2008; Panatier et al., 2011). Such spatial coupling could ultimately underlie the preferential occurrence of functional gliotransmission from astrocytic processes rather than from the soma (Gordon et al., 2009), entailing strict spatiotemporal requirements for Ca^{2+} signals to trigger release of gliotransmitters from the astrocyte and modulate synaptic transmission.

MODULATION OF SYNAPTIC RELEASE BY ASTROCYTIC GLUTAMATE AND ATP

Astrocyte-derived glutamate and ATP or adenosine can modulate synaptic transmission, either increasing neurotransmitter release (Araque et al., 1998a,b; Fiacco and McCarthy, 2004; Jourdain et al., 2007; Perea and Araque, 2007; Bonansco et al., 2011; Di Castro et al., 2011; Panatier et al., 2011) or decreasing it (Zhang et al., 2003, 2004a; Pascual et al., 2005; Andersson and Hansé, 2010, 2011) depending on the type of presynaptic receptor involved and the brain area (pathway **B** in **Figure 1B**; see also **Table 1**). At excitatory synapses in the hippocampal dentate gyrus, glutamate is released from surrounding astrocytic processes in close proximity to presynaptic NR_2B -containing NMDA receptors (**Figure 4A**). Activation of these receptors results in increased synaptic release and strengthening of synaptic transmission (Jourdain et al., 2007). At Schaffer collateral synapses in the CA1 hippocampal area a similar effect is mediated by presynaptic mGluRs (Fiacco and McCarthy, 2004; Perea and Araque, 2007). Besides directly targeting presynaptic receptors, astrocyte-released glutamate could also bind ionotropic receptors found along the axons of hippocampal CA3 pyramidal cells, broadening the profile of propagating action potentials (APs) (Sasaki et al., 2011). The broadened APs in turn, trigger larger Ca^{2+} elevations in presynaptic boutons, resulting in larger synaptic release probability.

Conversely, astrocyte-released ATP and its derivative adenosine, bind to presynaptic PRs, i.e. P_2Y_1 Rs or A_1 Rs, respectively, decreasing synaptic release (Zhang et al., 2003, 2004a; Pascual et al., 2005). However, astrocyte-derived adenosine, could also target $\text{A}_{2\text{A}}$ receptors which can increase synaptic release (Panatier et al., 2011) (omitted from **Figures 1A** and **4A** for simplicity). Both effects of adenosine—inhibitory via A_1 Rs and stimulatory via $\text{A}_{2\text{A}}$ Rs—have been described at hippocampal CA3-CA1 synapses (Zhang et al., 2003, 2004a; Pascual et al., 2005; Serrano et al., 2006; Panatier et al., 2011), and the prevalence of one on



the other likely depends on the level of synaptic activity (Panatier et al., 2011). This would be in line with observations at the frog's neuromuscular junction, where different stimulations of the tibial nerve differentially activated A₁Rs or A_{2A}Rs in association with different Ca²⁺ dynamics in the peri-junctional glial cell (Todd et al., 2010). Intriguingly, at hippocampal synapses different synaptic stimuli could differentially change the morphology of astrocyte perisynaptic processes (Haber et al., 2006; Lavielle et al., 2011) thus reshaping, in an activity-dependent fashion, the extracellular space of interaction between astrocyte-derived adenosine and presynaptic receptors. This could dynamically regulate access of adenosine to one receptor with respect to the other (Haber et al., 2006; Syková and Nicholson, 2008) ultimately modulating synaptic release in a complex fashion. The functional consequences of astrocytic remodeling were indeed demonstrated in the supraoptic nucleus, where dynamic changes in the astrocytic wrapping of synapses during lactation could regulate the extent of synaptic glutamate spillover and thereby control heterosynaptic depression of GABAergic transmission by presynaptic mGluRs (Oliet et al., 2001; Piet et al., 2004).

The opposite effects due to astrocytic glutamate or ATP/adenosine could endow astrocytes with the capacity to exert non-stereotyped bimodal control of synaptic transmission (Volterra and Meldolesi, 2005). On the other hand, the temporal concurrence of both these effects due to co-expression of inhibitory and stimulatory receptors at the same synaptic terminals (Shigemoto et al., 1997; Rebola et al., 2005), could result in occlusion, i.e. no net effect on synaptic release by the astrocyte (De Pittà et al., 2011). Alternatively, balanced activation (and possibly occlusion) of A₁ and A_{2A} receptors by astrocyte-derived adenosine could set synaptic release in basal conditions (Panatier et al., 2011). Hence, the ensuing regulation of synaptic transmission triggered by gliotransmitters in response to stimuli, could result instead from an unbalance of activation of these receptors rather than by the distinct activation of one receptor type with respect to the other, ultimately providing a high degree of complexity in the control of synaptic transmission by astrocytes. For the sake of clarity, in the following we will consider only release-decreasing effects of astrocyte ATP or its derivative, adenosine.

The details of the biochemical mechanism underlying modulation of synaptic release by astrocytic glutamate or ATP (or adenosine) likely depend on the type of targeted presynaptic receptors and are not fully understood (Pinheiro and Mulle, 2008). The simplest explanation would be that astrocytic glutamate and ATP lead to a modulation of presynaptic intracellular Ca²⁺ levels which eventually results in a modulation of synaptic release probability (Zucker and Regehr, 2002; Pinheiro and Mulle, 2008) with significant repercussions on synaptic plasticity, including short-term depression and facilitation.

Short-term facilitation and depression can be assessed by measuring the paired-pulse ratio (PPR), i.e. the ratio between the amplitudes of successive postsynaptic currents (PSCs) recorded in response to a pair of action potentials in rapid succession as illustrated in **Figure 4B** (Zucker and Regehr, 2002). When the value of the peak postsynaptic current associated to the second incoming spike is larger than the peak current recorded in coincidence with

the first spike (*green traces*), then synaptic release is increasing for incoming spikes, i.e. facilitation occurs, and the corresponding PPR is larger than unity. Conversely, when the second peak is less than the first peak (*magenta traces*), this marks a decrease of neurotransmitter release from the presynaptic terminal which reflects depression, and the corresponding value of PPR is less than one. It should be noted that, by varying the interval between two pulses, the same synapse can be either depressed or facilitated. For simplicity we omit the interpulse interval as a variable in this description.

In general, in basal conditions, i.e. in response to an individual action potential, the value of synaptic release probability of an individual synapse or of an ensemble of synapses defines the nature of synaptic transmission at that synapse/ensemble, namely whether it is facilitating or depressing, with low values of probability favoring facilitation and high values favoring depression (Abbott and Regehr, 2004) (**Figure 4B, right**). Thus, any modulation of synaptic release probability by gliotransmitters that changes the PPR from below unity to values above it or vice versa, could switch the mode of synaptic transmission from depressing to facilitating or vice versa. This scenario was theoretically addressed in De Pittà et al. (2011) and it was shown to substantially agree with experiments. Indeed, at hippocampal synapses, the increase of synaptic release probability due to astrocytic glutamate correlates with a *decrease* of the PPR (Jourdain et al., 2007; Perea and Araque, 2007; Bonansco et al., 2011). Conversely, a decrease of synaptic release due to the action of astrocyte-derived ATP (or adenosine) is accompanied by an *increase* of the PPR (Zhang et al., 2003, 2004a). The frequency of astrocytic glutamate (Glu) and/or ATP exocytosis are likely crucial in the regulation of the extent of the modulation of synaptic release by astrocytes (De Pittà et al., 2011), a principle illustrated in **Figure 4C**. This figure shows the simulated peak postsynaptic currents for the same synapse in response to 100 trials of presynaptic spike trains with identical statistics (*raster plots*). Each column in **Figure 4C** corresponds to a different frequency of astrocytic Ca²⁺ pulses (*top row*), yielding to different frequencies of gliotransmitter release when Ca²⁺ crosses the threshold for exocytosis (*dashed red line*) (see also **Figure 3B**). Colors in the raster plots refer to paired-pulse plasticity quantified by PPR. For two consecutive presynaptic spikes, if the second spike releases synaptic neurotransmitter more than the first spike, then it is PPR > 1, paired-pulse facilitation is observed, and the peak postsynaptic current associated to the second spike is colored in *green*. Vice versa, if the amount of neurotransmitter released by the second spike is less than that released by the first one, then PPR < 1, that is paired-pulse depression occurs, and the second peak postsynaptic current is colored in *magenta*.

In absence of astrocytic gliotransmitter (*left column*), the raster plot of a facilitating synapse (*middle row*) shows predominant paired-pulse facilitation, i.e. mostly *green* dots. However, in presence of release of glutamate from the astrocyte (+Glu), as explained above, the stimulatory effect of this gliotransmitter on synaptic release changes paired-pulse plasticity which is marked by the appearance of *magenta* bands in the raster plots (*middle and right columns*). These bands locate time intervals where paired-pulse depression becomes prominent (i.e. predominance

of *magenta* dots) on the overall paired-pulse facilitation background (*green dots*). Notably these bands are almost in coincidence with glutamate release from the astrocyte (*top row*) and their number increases with the glutamate release frequency. Therefore, for the same time window, the same originally-facilitating synapse gets increasingly depressing as the rate of glutamate release from the astrocyte increases. The opposite is observed for an otherwise depressing synapse (i.e. predominance of paired-pulse depression, thus mostly *magenta* dots in the *bottom left* raster plot). In this case, in presence of release of ATP and its derivative adenosine (+ATP/Adn) from the astrocyte, due to the inhibitory effect of astrocyte-derived purines on synaptic release considered in this example, *green* bands appear in the raster plots which mark the onset of periods of predominant paired-pulse facilitation. The number of these bands grows for higher rates of ATP release from the astrocyte so that the same originally-depressing synapse behaves more akin of a facilitating one as the rate of ATP release from the astrocyte increases.

An intriguing prediction that follows from the above arguments is that the frequency of gliotransmitter release, by modulating synaptic release probability at basal conditions, could dynamically control the nature of synaptic transmission as elucidated in **Figure 5A**. In particular, under certain conditions, a threshold frequency for gliotransmitter release (*blue line*) could exist above which the astrocyte can switch the nature of synaptic transmission, turning depressing synapses into facilitating or vice versa, facilitating synapses into depressing (De Pittà et al., 2011). Hence, the plasticity mode at such synapses is not fixed but rather is set by the release rate of gliotransmitters from neighboring astrocytic processes.

Remarkably, the threshold frequency of gliotransmitter release that discriminates between facilitation and depression at one synapse can be as low as 0.05 Hz or less (as in the case of **Figure 5A**) thus falling within the range of Ca^{2+} oscillations observed in astrocytes in basal conditions (Parri et al., 2001; Aguado et al., 2002; Bonansco et al., 2011; Di Castro et al., 2011). In such conditions as shown in **Figure 5B** (*top*), intracellular Ca^{2+} levels in astrocytes spontaneously fluctuate in a highly stochastic fashion (Hirase et al., 2004; Di Castro et al., 2011) and can similarly cross the threshold for gliotransmitter release. The ensuing release of gliotransmitter, however, could be sufficient to tonically set the basal tone of synaptic transmission (**Figure 5B**). Inasmuch as the rate of gliotransmitter release could also correlate with the incoming synaptic stimulus through Ca^{2+} dynamics in the astrocytes (Aguado et al., 2002; Di Castro et al., 2011), this scenario discloses the possibility that astrocytes act as endogenous regulators of the efficacy of synaptic transmission (Haydon, 2001; Giaume et al., 2010; Halassa and Haydon, 2010; Di Castro et al., 2011; Panatier et al., 2011). That is, through integration of synaptic activity by means of their Ca^{2+} signals, astrocytes could adapt synaptic strength according to the history of the synapse.

IMPLICATIONS OF ASTROCYTE MODULATION OF SYNAPTIC TRANSMISSION AND PLASTICITY

The effect of modulation of synaptic release probability by astrocytic gliotransmitters may decay more slowly than the Ca^{2+}

elevation triggering astrocyte exocytosis (Fiacco and McCarthy, 2004; Serrano et al., 2006), and still be present upon gliotransmitter release by following Ca^{2+} increases (Volman et al., 2007). Therefore astrocytic Ca^{2+} activity resulting in high rates of gliotransmitter release, could bring forth long-lasting modulations of synaptic release. Insofar as synaptically-released neurotransmitter could shape postsynaptic signaling at the basis of long-term synaptic plasticity (Bliss and Collingridge, 1993), persistent modulations of synaptic release probability by astrocyte could ultimately contribute to long-term changes of synaptic strength underlying neural processing, memory formation and storage of information. Support to this scenario comes from studies on hippocampal synapses for which the temporal coincidence of postsynaptic depolarization with the increase of Ca^{2+} elevations in neighboring astrocytes was shown to induce long-term potentiation (LTP) of synaptic transmission (Perea and Araque, 2007; Navarrete and Araque, 2010; Navarrete et al., 2012a,b). Notably, this form of LTP is independent of the activation of postsynaptic NMDARs but rather, it depends on glutamate released from astrocytes, which persistently potentiates synaptic transmitter release through activation of presynaptic mGluRs (Perea and Araque, 2007). On the other hand, at synapses between excitatory neurons in layers 4 and 2/3 in the rat barrel cortex, activity-dependent induction of long-term depression (LTD) also requires astrocyte Ca^{2+} signaling (Min and Nevian, 2012). At these synapses, postsynaptically-released endocannabinoids mediate Ca^{2+} -dependent release of glutamate from astrocytes which targets presynaptic NMDARs bringing forth LTD (Sjöström et al., 2003; Rodríguez-Moreno and Paulsen, 2008).

The differential induction of LTP and LTD in neighboring synapses has been suggested to determine the size and shape of cortical functional topographic units such as ocular dominance columns in the primary visual cortex and whisker barrels in the primary somatosensory cortex (Feldman and Brecht, 2005; Hensch, 2005; Li et al., 2009). Thus, the possible involvement of astrocytes in LTP and LTD suggests that these cells could contribute to the plasticity of cortical maps and the development of corresponding sensory representations (Rossi, 2012).

Modulation of synaptic release probability by astrocytic gliotransmitters could also alter the temporal order of correlated pre- and postsynaptic spiking that critically dictates spike-timing-dependent plasticity (STDP) (Pascual et al., 2005). Inhibition of spontaneous glutamate release from astrocytes in hippocampal CA1 synapses, consistent with a decrease of synaptic release probability, was indeed reported to modify the threshold for induction of spike-timing-dependent LTP (Bonansco et al., 2011). In this fashion, astrocytes by gliotransmission-mediated regulations of synaptic release probability could control not only different mechanisms of synaptic plasticity but also the threshold of synaptic activity required for their onset, thus playing a role in metaplasticity too, that is the plasticity of synaptic plasticity (Abraham, 2008).

Gliotransmitters do not control synaptic plasticity only via presynaptic actions, but also by actions on postsynaptic receptors. Indeed, the induction of LTP itself appears to be uniquely




FIGURE 5 | Short-term synaptic plasticity is physiologically set by the rate of gliotransmitter release from the astrocyte. (A) The rate of glutamate or ATP release from the astrocyte could differently affect basal synaptic release probability. In particular, a threshold frequency for the release of these gliotransmitters could exist (*blue vertical line*) beyond which a depressing synapse could turn into facilitating (*right, green area*) or vice versa, a facilitating synapse could become depressing (*left, magenta area*). Adapted from De Pittà et al. (2011). **(B)** In basal conditions,

synaptic release is due to sporadic neuronal network firing and the possible frequency of Ca^{2+} fluctuations beyond the threshold for exocytosis (*dashed red line*) in the astrocyte (*top trace*). In this fashion plastic changes in paired-pulse ratio could be inherently regulated by astrocytic gliotransmitters, as shown here for the case of an originally facilitating synapse under the effect of astrocytic glutamate exocytosis (20 trials with identical input statistics). Adapted from Berry et al. (2011). The color code for the raster plot is the same as in **Figure 4C**.

controlled by astrocytes through the release of D-serine (Santello and Volterra, 2010) (pathway C in **Figure 1B**). Both at hippocampal and cortical synapses astrocytic D-serine rather than glycine, is the endogenous co-agonist of synaptic NMDA receptors (Henneberger et al., 2010; Takata et al., 2011; Papouin et al., 2012). By controlling the level of co-agonist site occupancy of postsynaptic NMDARs, astrocytic D-serine affects the level of activation of these receptors and thus activity-dependent long-term synaptic changes (Bains and Oliet, 2007). In particular, the additional burst of activation of postsynaptic NMDARs (about 25% more) induced by astrocyte D-serine release seems necessary for LTP induction (Henneberger et al., 2010). Moreover, in conditions of reduced synaptic coverage by astrocytes, such as during lactation in the hypothalamus, higher presynaptic activity is required to obtain LTP while the same level of activity that normally induces LTP results instead in LTD (Panatier et al., 2006). In such conditions, astrocyte-released D-serine is diluted in the larger extracellular space resulting in a reduced number of postsynaptic NMDARs recruited by synaptic activity, which ultimately translates into smaller postsynaptic Ca^{2+} increases.

Therefore, experimental protocols that would be expected to cause LTP, elicit LTD instead. This is in agreement with the Bienenstock-Cooper-Munro (BCM) model for variation of the threshold for LTP, which predicts that the relationship between synaptic activity and persistent changes in synaptic strength can vary according to the number of NMDARs available during synaptic activation (Bienenstock et al., 1982; Abraham and Bear, 1996). Effectively, by adjusting the D-serine occupancy of the NMDAR co-agonist-binding site, astrocytes can shift the relationship between activity and synaptic strength (Panatier et al., 2006).

Besides D-serine, experimental evidence hints that glutamate and ATP released from astrocytes could also directly bind postsynaptically-located receptors and accordingly, play a role in regulation of long-term synaptic plasticity. In the paraventricular nucleus of the hypothalamus for example, ATP released from astrocytes could directly target postsynaptic P_2X_7 receptors, promoting insertion of postsynaptic AMPARs which results in LTP of synaptic transmission (Gordon et al., 2005). Interestingly, ATP release is mediated by Ca^{2+} dynamics

triggered in astrocytes by noradrenergic afferents which, in the hypothalamus, generally lack direct postsynaptic contacts (Sawyer and Clifton, 1980), thus hinting that signaling in this vital homeostatic circuit may require dynamic neuron-glia interactions.

Glutamate released from astrocyte could also target extrasynaptically-located NR₂B-containing NMDA receptors at postsynaptic terminals, triggering slow inward currents (SICs) (Fellin et al., 2004; D'Ascenzo et al., 2007; Navarrete and Araque, 2008; Bardoni et al., 2010; Pirttimaki et al., 2011; Navarrete et al., 2012a,b) mainly mediated by Ca²⁺ ions (Cull-Candy et al., 2001), whose depolarizing action could affect postsynaptic neuronal firing (D'Ascenzo et al., 2007; Pirttimaki et al., 2011). In the primary visual cortex, nucleus basalis-mediated cholinergic activation of astrocytes mediates an increase of SICs frequency which correlates with a long-lasting increase of firing activity in visual responses of V1 excitatory neurons (Chen et al., 2012). The ensuing modulations of firing activity of these neurons by astrocyte-mediated SICs might ultimately affect STDP at individual synapses controlling orientation-specific responses of V1 neurons to visual stimuli (Jia et al., 2010).

Modulation of synaptic transmission by Ca²⁺-dependent gliotransmission may not be limited to the very synapses that trigger Ca²⁺ activity in the astrocyte but it could also affect farther synaptic domains in a multimodal fashion (Kozlov et al., 2006), depending both on the morphology of the sites of astrocyte-synapse reciprocal communication (Ventura and Harris, 1999; Haber et al., 2006) and the functional connectivity of the astrocytic network (Pannasch et al., 2011). High-frequency activity of a Schaffer collateral fiber can trigger the potentiation of synaptic transmission at the same fiber but heterosynaptic suppression of another, adjacent fiber, by inducing ATP release from an astrocyte interposed between the two fibers (Zhang et al., 2003, 2004a; Pascual et al., 2005). In the somatosensory cortex in particular, astrocyte-mediated heterosynaptic suppression could modulate GABAergic inhibitory transmission (Benedetti et al., 2011) which plays a dominant role in the control of cortical neuronal excitability (Swadlow, 2002). Given that both experimental observations and theoretical arguments suggest that excitation and inhibition are globally balanced in cortical circuits (Shadlen and Newsome, 1994; Troyer and Miller, 1997; Shu et al., 2003; Haider et al., 2006), one may speculate that this mechanism could be involved in gating of signal transmission (Buzsáki, 2010). That is, by modulating inhibitory synaptic transmission, astrocytes could favor network excitation resulting in neuronal firing consistent with the transmission, i.e. gating “on,” of specific stimuli rather than others (Vogels and Abbott, 2009).

The latter idea could also bring to a possible role of astrocyte signaling in coherent function of neural networks underlying potential behavioral states (Engel et al., 2001). In cortical slices, for example, stimulation of a single astrocyte was reported to activate large portions of the astrocytic network and to result in an increase of synchronized neuronal depolarizations (Poskanzer and Yuste, 2011). This phenomenon was suggested

to modulate the induction of cortical UP and DOWN states, possibly involved in determining the oscillatory activity observed in slow-wave sleep (Fellin et al., 2004; Halassa and Haydon, 2010; Poskanzer and Yuste, 2011), and is consistent with reports of sleep perturbations in mice lacking astrocytic gliotransmitter exocytosis (Fellin et al., 2009; Halassa et al., 2009; Fellin et al., 2012).

CONCLUSIONS

A large body of evidence has accumulated over the last years, suggesting an active role of astrocytes in many brain functions. Collectively these data have fuelled the concept that synapses could be tripartite rather than bipartite, since in addition to the pre- and post-synaptic terminals, the astrocyte could be an active element in synaptic transmission (Araque et al., 1999; Haydon, 2001). While the tripartite synapse concept captures well the essence of astrocyte-regulated synapses, the inclusion of astrocytic signaling within our current knowledge of synaptic transmission could add more than just one level of complexity. Existing evidence suggests that astrocytes could produce not just tonic and diffuse modulatory influences on synapses but also engage in more focused, spatially precise and constrained communications with synaptic terminals (Anderson and Swanson, 2000; Jourdain et al., 2007; Santello and Volterra, 2009; Bergersen et al., 2012). This calls to rethink the definition of a functional synapse, to include the contribution from surrounding astrocytes. To conclude, the growing appreciation that astrocytes can regulate synaptic information at many levels, from activity of single synapses to network levels and behavioral states (Fellin et al., 2009; Halassa et al., 2009; Zorec et al., 2012) changes our understanding of brain communication and the role of glial cells in synaptic transmission. This resulting novel scenario offers an enticing platform for future theoretical investigations that we are just beginning to appreciate in its potential far-reaching implications.

ACKNOWLEDGMENTS

The authors would like to thank Richard Robitaille, Mirko Santello, and Ari Barzilai for insightful conversations and Nicolas Liaudet for critical reading of the manuscript. The authors' research was supported by the Tauber Family Foundation (Maurizio De Pittà and Eshel Ben-Jacob); the Italy-Israel Joint Neuroscience Lab (Maurizio De Pittà); the Maguy-Glass Chair in Physics of Complex Systems at Tel Aviv University (Eshel Ben-Jacob); the Center for Theoretical Biological Physics at Rice University sponsored by the National Science Foundation (NSF) (grant PHY-0822283) (Eshel Ben-Jacob); the NSF program in Physics of Living Systems (PoLS) at the University of Michigan (grant PHY-1058034) (Eshel Ben-Jacob); the National Institute of Health (grants R01 NS059740 and R01 NS060870 to T. Sejnowski and M. Bazhenov) (Vladislav Volman); the High Council for Scientific and Technological Cooperation between France-Israel (Hugues Berry); the National Science Foundation (grant CBET 0943343) (Vladimir Parpura); and the Swiss National Science Foundation (grant 31003A-140999 and NCCRs “Synapsy” and “Transcure”) (Andrea Volterra).

REFERENCES

- Abbott, L. F., and Regehr, W. G. (2004). Synaptic computation. *Nature* 431, 796–803.
- Abbracchio, M. P., Burnstock, G., Verkhratsky, A., and Zimmermann, H. (2009). Purinergic signalling in the nervous system: an overview. *Trends Neurosci.* 32, 19–29.
- Abraham, W. (2008). Metaplasticity: tuning synapses and networks for plasticity. *Nat. Rev. Neurosci.* 9, 387–387.
- Abraham, W., and Bear, M. (1996). Metaplasticity: the plasticity of synaptic plasticity. *Trends Neurosci.* 19, 126–130.
- Aguado, F., Espinosa-Parrilla, J. F., Carmona, M. A., and Soriano, E. (2002). Neuronal activity regulates correlated network properties of spontaneous calcium transients in astrocytes *in situ*. *J. Neurosci.* 22, 9430–9444.
- Agulhon, C., Fiacco, T. A., and McCarthy, K. D. (2010). Hippocampal short- and long-term plasticity are not modulated by astrocyte Ca^{2+} signalling. *Science* 327, 1250–1254.
- Agulhon, C., Petravic, J., McMullen, A. B., Sweger, E. J., Minton, S. K., Taves, S. R., et al. (2008). What is the role of astrocyte calcium in neurophysiology? *Neuron* 59, 932–946.
- Allbritton, N. L., Meyer, T., and Stryer, L. (1992). Range of messenger action of calcium ion and inositol 1,4,5-trisphosphate. *Science* 260, 107–260. Abstract.
- Andersson, M., and Hanse, E. (2010). Astrocytes impose postburst depression of release probability at hippocampal glutamate synapses. *J. Neurosci.* 30, 5776–5780.
- Anderson, M. C., and Swanson, R. A. (2000). Astrocyte glutamate transport: review of properties, regulation, and physiological functions. *Glia* 32, 1–14.
- Andersson, M. S., and Hanse, E. (2011). Astrocyte-mediated short-term synaptic depression in the rat hippocampal CA1 area: two modes of decreasing release probability. *BMC Neurosci.* 12:87. doi: 10.1186/1471-2202-12-87
- Araque, A., Martín, E., Perea, G., Arellano, J., and Buño, W. (2002). Synaptically released acetylcholine evokes Ca^{2+} elevations in astrocytes in hippocampal slices. *J. Neurosci.* 22, 2443–2450.
- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1998a). Glutamate-dependent astrocyte modulation of synaptic transmission between cultured hippocampal neurons. *Eur. J. Neurosci.* 10, 2129–2142.
- Araque, A., Sanzgiri, R. P., Parpura, V., and Haydon, P. G. (1998b). Calcium elevation in astrocytes causes an NMDA receptor-dependent increase in the frequency of miniature synaptic currents in cultured hippocampal neurons. *J. Neurosci.* 18, 6822–6829.
- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1999). Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci.* 23, 208–215.
- Arizono, M., Bannai, H., Nakamura, K., Niwa, F., Enomoto, M., Matsuura, T., et al. (2012). Receptor-selective diffusion barrier enhances sensitivity of astrocytic processes to metabotropic glutamate receptor stimulation. *Sci. Signal.* 5, ra27.
- Auld, D., and Robitaille, R. (2003). Glial cells and neurotransmission: an inclusive view of synaptic function. *Neuron* 40, 389–400.
- Bains, J. S., and Oliet, S. H. R. (2007). Glia: they make your memories stick! *Trends Neurosci.* 30, 417–424.
- Balosso, S., Ravizza, T., Pierucci, M., Calcagno, E., Invernizzi, R., Di Giovanni, G., et al. (2009). Molecular and functional interactions between tumor necrosis factor- α receptors and the glutamatergic system in the mouse hippocampus: implications for seizure susceptibility. *Neuroscience* 161, 293–300.
- Bardoni, R., Ghirri, A., Zonta, M., Betelli, C., Vitale, G., Ruggieri, V., et al. (2010). Glutamate-mediated astrocyte-to-neuron signalling in the rat dorsal horn. *J. Physiol.* 588, 831–846.
- Barlow, H. (1996). Intraneuronal information processing, directional selectivity and memory for spatio-temporal sequences. *Network* 7, 251–259.
- Barres, B. (2008). The mystery and magic of glia: a perspective on their roles in health and disease. *Neuron* 60, 430–440.
- Beattie, E., Stellwagen, D., Morishita, W., Bresnahan, J., Ha, B., von Zastrow, M., et al. (2002). Control of synaptic strength by glial TNF α . *Science* 295, 2282–2285.
- Beierlein, M., and Regehr, W. (2006). Brief bursts of parallel fiber activity trigger calcium signals in Bergmann glia. *J. Neurosci.* 26, 6958–6967.
- Bekar, L., He, W., and Nedergaard, M. (2008). Locus coeruleus α -adrenergic-mediated coactivation of cortical astrocytes *in vivo*. *Cereb. Cortex* 18, 2789–2795.
- Bélaïr, E., Vallée, J., and Robitaille, R. (2010). *In vivo* long-term synaptic plasticity of glial cells. *J. Physiol.* 588, 1039–1056.
- Benedetti, B., Matyash, V., and Kettenmann, H. (2011). Astrocytes control GABAergic inhibition of neurons in the mouse barrel cortex. *J. Physiol.* 589, 1159–1172.
- Bennett, M., Farnell, L., and Gibson, W. (2008). A quantitative model of cortical spreading depression due to purinergic and gap-junction transmission in astrocyte networks. *Biophys. J.* 95, 5648–5660.
- Bergersen, L. H., and Gundersen, V. (2009). Morphological evidence for vesicular glutamate release from astrocytes. *Neuroscience* 158, 260–265.
- Bergersen, L. H., Morland, C., Ormel, L., Rinholm, J. E., Larsson, M., Wold, J. F. H., et al. (2012). Immunogold detection of L-glutamate and D-serine in small synaptic-like microvesicles in adult hippocampal astrocytes. *Cereb. Cortex* 22, 1690–1697.
- Bernardinelli, Y., Salmon, C., Jones, E., Farmer, W., Stellwagen, D., and Murai, K. (2011). Astrocytes display complex and localized calcium responses to single-neuron stimulation in the hippocampus. *J. Neurosci.* 31, 8905–8919.
- Berridge, M. J. (1997). The AM and FM of calcium signaling. *Nature* 389, 759–760.
- Berridge, M. J., Bootman, M. D., and Roderick, H. L. (2003). Calcium signalling: dynamics, homeostasis and remodelling. *Nat. Rev.* 4, 517–529.
- Berry, H., De Pittà, M., Volman, V., and Ben-Jacob, E. (2011). “Astrocyte regulation of presynaptic plasticity,” in *2011 Neuroscience Meeting Planner* (Washington, DC: Society for Neuroscience) [Online].
- Bezprozvanny, I., Watras, J., and Ehrlich, B. E. (1991). Bell-shaped calcium-response curves of $Ins(1,4,5)P_3$ - and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature* 351, 751–754.
- Bezzi, P., Carmignoto, G., Pasti, L., Vesce, S., Rossi, D., Lodi Rizzini, B., et al. (1998). Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. *Nature* 391, 281–285.
- Bezzi, P., Domercq, M., Brambilla, L., Galli, R., Schols, D., De Clercq, E., et al. (2001). CXCR4-activated astrocyte glutamate release via TNF α : amplification by microglia triggers neurotoxicity. *Nat. Neurosci.* 4, 702–710.
- Bezzi, P., Gundersen, V., Galbete, J. L., Seifert, G., Steinhäuser, C., Pilati, E., et al. (2004). Astrocytes contain a vesicular compartment that is competent for regulated exocytosis of glutamate. *Nat. Neurosci.* 7, 613–620.
- Bienenstock, E., Cooper, L., and Munro, P. (1982). Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *J. Neurosci.* 2, 32–48.
- Blaustein, M., and Golovina, V. (2001). Structural complexity and functional diversity of endoplasmic reticulum Ca^{2+} stores. *Trends Neurosci.* 24, 602–608.
- Bliss, T., and Collingridge, G. (1993). A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361, 31–39.
- Blomstrand, F., and Giaume, C. (2006). Kinetics of endothelin-induced inhibition and glucose permeability of astrocyte gap junctions. *J. Neurosci. Res.* 83, 996–1003.
- Bonansco, C., Couve, A., Perea, G., Ferradas, C. A., Roncagliolo, M., and Fuenzalida, M. (2011). Glutamate released spontaneously from astrocytes sets the threshold for synaptic plasticity. *Eur. J. Neurosci.* 33, 1483–1492.
- Bootman, M., Lipp, P., and Berridge, M. (2001). The organisation and functions of local Ca^{2+} signals. *J. Cell Sci.* 114, 2213–2222.
- Bowser, D., and Khakh, B. (2004). ATP excites interneurons and astrocytes to increase synaptic inhibition in neuronal networks. *J. Neurosci.* 24, 8606–8620.
- Bowser, D. N., and Khakh, B. S. (2007). Two forms of single-vesicle astrocyte exocytosis imaged with total internal reflection fluorescence microscopy. *Proc. Natl. Acad. Sci. U.S.A.* 104, 4212–4217.
- Bushong, E. A., Martone, M. E., Jones, Y. Z., and Ellisman, M. H. (2002). Protoplasmic astrocyte in CA1 stratum radiatum occupy separate anatomical domains. *J. Neurosci.* 22, 183–192.
- Buzsáki, G. (2010). Neural syntax: cell assemblies, synapses, and readers. *Neuron* 68, 362–385.
- Carmignoto, G. (2000). Reciprocal communication systems between astrocytes and neurones. *Prog. Neurobiol.* 62, 561–581.
- Chan-Ling, T., and Stone, J. (1991). Factors determining the morphology and distribution of astrocytes in the cat retina: a ‘contact-spacing’ model of astrocyte interaction. *J. Comp. Neurol.* 303, 387–399.
- Chen, N., Sugihara, H., Sharma, J., Perea, G., Petravic, J., Le, C., et al.

- (2012). Nucleus basalis enabled stimulus specific plasticity in the visual cortex is mediated by astrocytes. *Proc. Natl. Acad. Sci. U.S.A.* 109, E2832–E2841.
- Chen, X., Wang, L., Zhou, Y., Zheng, L.-H., and Zhou, Z. (2005). “Kiss-and-run” glutamate secretion in cultured and freshly isolated rat hippocampal astrocytes. *J. Neurosci.* 25, 9236–9243.
- Coco, S., Calegari, F., Pravettoni, E., Pozzi, D., Taverna, E., Rosa, P., et al. (2003). Storage and release of ATP from astrocytes in culture. *J. Biol. Chem.* 278, 1354–1362.
- Codazzi, F., Teruel, M. N., and Meyer, T. (2001). Control of astrocyte Ca^{2+} oscillations and waves by oscillating translocation and activation of protein kinase C. *Curr. Biol.* 11, 1089–1097.
- Communi, D., Gevaert, K., Demol, H., Vandekerckhove, J., and Erneux, C. (2001). A novel receptor-mediated regulation mechanism of type I inositol polyphosphate 5-phosphatase by calcium/calmodulin-dependent protein kinase II phosphorylation. *J. Biol. Chem.* 276, 38738–38747.
- Communi, D., Vanweyenberg, V., and Erneux, C. (1997). d-myo-inositol 1,4,5-trisphosphate 3-kinase A is activated by receptor activation through a calcium: calmodulin-dependent protein kinase II phosphorylation mechanism. *EMBO J.* 16, 1943–1952.
- Crippa, D., Schenk, U., Francolini, M., Rosa, P., Verderio, C., Zonta, M., et al. (2006). Synaptobrevin2-expressing vesicles in rat astrocytes: insights into molecular characterization, dynamics and exocytosis. *J. Physiol.* 570, 567–582.
- Cull-Candy, S., Brickley, S., and Farrant, M. (2001). NMDA receptor subunits: diversity, development and disease. *Curr. Opin. Neurobiol.* 11, 327–335.
- D’Ascenzo, M., Fellin, T., Terunuma, M., Revilla-Sanchez, R., Meaney, D., Auberson, Y., et al. (2007). mGluR5 stimulates gliotransmission in the nucleus accumbens. *Proc. Natl. Acad. Sci. U.S.A.* 104, 1995.
- Del Castillo, J., and Katz, B. (1954). Quantal components of the endplate potential. *J. Physiol.* 124, 560–573.
- Deng, Q., Terunuma, M., Fellin, T., Moss, S. J., and Haydon, P. (2011). Astrocytic activation of A1 receptors regulates the surface expression of NMDA receptors through a src kinase dependent pathway. *Glia* 59, 1084–1093.
- De Pittà, M., Volman, V., Levine, H., and Ben-Jacob, E. (2009a). Multimodal encoding in a simplified model of intracellular calcium signaling. *Cogn. Processing* 10(Suppl. 1), 55–70.
- De Pittà, M., Goldberg, M., Volman, V., Berry, H., and Ben-Jacob, E. (2009b). Glutamate-dependent intracellular calcium and IP_3 oscillating and pulsating dynamics in astrocytes. *J. Biol. Phys.* 35, 383–411.
- De Pittà, M., Volman, V., Berry, H., and Ben-Jacob, E. (2011). A tale of two stories: astrocyte regulation of synaptic depression and facilitation. *PLoS Comput. Biol.* 7:e1002293. doi: 10.1371/journal.pcbi.1002293
- De Pittà, M., Volman, V., Levine, H., Pioggia, G., De Rossi, D., and Ben-Jacob, E. (2008). Coexistence of amplitude and frequency modulations in intracellular calcium dynamics. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* 77, 030903(R).
- De Saint Jan, D., and Westbrook, G. (2005). Detecting activity in olfactory bulb glomeruli with astrocyte recording. *J. Neurosci.* 25, 2917–2924.
- Di Castro, M., Chuquet, J., Liaudet, N., Bhaukaurally, K., Santello, M., Bouvier, D., et al. (2011). Local Ca^{2+} detection and modulation of synaptic release by astrocytes. *Nat. Neurosci.* 14, 1276–1284.
- Domercq, M., Brambilla, L., Pilati, E., Marchaland, J., Volterra, A., and Bezzi, P. (2006). P2Y1 receptor-evoked glutamate exocytosis from astrocytes: control by tumor necrosis factor- α and prostaglandins. *J. Biol. Chem.* 281, 30684–30696.
- Dundwiddie, T. V., and Masino, S. A. (2001). The role and regulation of adenosine in the central nervous system. *Ann. Rev. Neurosci.* 24, 31–55.
- Dupont, G., Lokenye, E. F., and Challiss, R. A. (2011). A model for Ca^{2+} oscillations stimulated by the type 5 metabotropic glutamate receptor: an unusual mechanism based on repetitive, reversible phosphorylation of the receptor. *Biochimie* 93, 2132–2138.
- Emptage, N. J., Reid, C. A., and Fine, A. (2001). Calcium stores in hippocampal synaptic boutons mediate short-term plasticity, store-operated Ca^{2+} entry, and spontaneous transmitter release. *Neuron* 29, 197–208.
- Engel, A., Fries, P., and Singer, W. (2001). Dynamic predictions: oscillations and synchrony in top-down processing. *Nat. Rev. Neurosci.* 2, 704–716.
- Enkvist, M. O., and McCarthy, K. D. (1992). Activation of protein kinase C blocks astroglial gap junction communication and inhibits the spread of calcium waves. *J. Neurochem.* 59, 519–526.
- Ermentrout, G., and Terman, D. (2010). *Mathematical Foundations of Neuroscience*. New York, NY: Springer Verlag.
- Falcke, M. (2004). Reading the patterns in living cells – the physics of Ca^{2+} signaling. *Adv. Phys.* 53, 255–440.
- Feldman, D., and Brecht, M. (2005). Map plasticity in somatosensory cortex. *Science* 310, 810.
- Fellin, T., Ellenbogen, J. M., De Pittà, M., Ben-Jacob, E., and Halassa, M. (2012). Astrocyte regulation of sleep circuits: experimental and modeling perspectives. *Front. Comput. Neurosci.* 6:65. doi: 10.3389/fncom.2012.00065
- Fellin, T., Halassa, M., Terunuma, M., Succol, F., Takano, H., Frank, M., et al. (2009). Endogenous nonneuronal modulators of synaptic transmission control cortical slow oscillations *in vivo*. *Proc. Natl. Acad. Sci. U.S.A.* 106, 15037–15042.
- Fellin, T., Pascual, O., Gobbo, S., Pozzan, T., Haydon, P. G., and Carmignoto, G. (2004). Neuronal synchrony mediated by astrocytic glutamate through activation of extrasynaptic NMDA receptors. *Neuron* 43, 729–743.
- Ferguson, A., Christensen, R., Gensel, J., Miller, B., Sun, F., Beattie, E., et al. (2008). Cell death after spinal cord injury is exacerbated by rapid TNF α -induced trafficking of GluR2-lacking AMPARs to the plasma membrane. *J. Neurosci.* 28, 11391–11400.
- Fiacco, T. A., Agulhon, C., Taves, S. R., Petravic, J., Casper, K. B., Dong, X., et al. (2007). Selective stimulation of astrocyte calcium *in situ* does not affect neuronal excitatory synaptic activity. *Neuron* 54, 611–626.
- Fiacco, T. A., and McCarthy, K. D. (2004). Intracellular astrocyte calcium waves *in situ* increase the frequency of spontaneous AMPA receptor currents in CA1 pyramidal neurons. *J. Neurosci.* 24, 722–732.
- Fukaya, M., Uchigashima, M., Nomura, S., Hasegawa, Y., Kikuchi, H., and Watanabe, M. (2008). Predominant expression of phospholipase C β 1 in telencephalic principal neurons and cerebellar interneurons, and its close association with related signaling molecules in somatodendritic neuronal elements. *Eur. J. Neurosci.* 28, 1744–1759.
- García-Marqués, J., and López-Mascaraque, L. (2012). Clonal identity determines astrocyte cortical heterogeneity. *Cereb. Cortex*. doi: 10.1093/cercor/bhs134. [Epub ahead of print].
- Giaume, C., Koulakoff, A., Roux, L., Holcman, D., and Rouach, N. (2010). Astroglial networks: a step further in neuroglial and gliovascular interactions. *Nat. Rev. Neurosci.* 11, 87–99.
- Goldberg, M., De Pittà, M., Volman, V., Berry, H., and Ben-Jacob, E. (2010). Nonlinear gap junctions enable long-distance propagation of pulsating calcium waves in astrocyte networks. *PLoS Comput. Biol.* 6:e1000909. doi: 10.1371/journal.pcbi.1000909
- Gordon, G., Baimoukhametova, D., Hewitt, S., Rajapaksha, W., Fisher, T., and Bains, J. (2005). Norepinephrine triggers release of glial ATP to increase postsynaptic efficacy. *Nat. Neurosci.* 8, 1078–1086.
- Gordon, G. R. J., Iremonger, K. J., Kantevari, S., Ellis-Davies, G. C. R., MacVicar, B. A., and Bains, J. S. (2009). Astrocyte-mediated distributed plasticity at hypothalamic glutamate synapses. *Neuron* 64, 391–403.
- Grosche, J., Matyash, V., Möller, T., Verkhratsky, A., Reichenbach, A., and Kettenmann, H. (1999). Microdomains for neuron-glia interaction: parallel fiber signaling to bergmann glial cells. *Nature* 2, 139–143.
- Guthrie, P. B., Knappenberger, J., Segal, M., Bennett, M. V., Charles, A. C., and Kater, S. B. (1999). ATP released from astrocytes mediates glial calcium waves. *J. Neurosci.* 19, 520–528.
- Haas, B., Schipke, C. G., Peters, O., Sohl, G., Willecke, K., and Kettenmann, H. (2006). Activity-dependent ATP-waves in the mouse neocortex are independent from astrocytic calcium waves. *Cereb. Cortex* 16, 237–246.
- Haber, M., Zhou, L., and Murai, K. K. (2006). Cooperative astrocyte and dendritic spine dynamics at hippocampal excitatory synapses. *J. Neurosci.* 26, 8881–8891.
- Haider, B., Duque, A., Hasenstaub, A., and McCormick, D. (2006). Neocortical network activity *in vivo* is generated through a dynamic balance of excitation and inhibition. *J. Neurosci.* 26, 4535–4545.
- Halassa, M., Florian, C., Fellin, T., Munoz, J., Lee, S., Abel, T., et al. (2009). Astrocytic modulation of sleep homeostasis and cognitive

- consequences of sleep loss. *Neuron* 61, 213–219.
- Halassa, M., and Haydon, P. (2010). Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annu. Rev. Physiol.* 72, 335–355.
- Haydon, P. G. (2001). Glia: listening and talking to the synapse. *Nat. Rev. Neurosci.* 2, 185–193.
- Haydon, P. G., and Carmignoto, G. (2006). Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol. Rev.* 86, 1009–1031.
- Henneberger, C., Papouin, T., Oliet, S. H. R., and Rusakov, D. A. (2010). Long-term potentiation depends on release of D-serine from astrocytes. *Nature* 463, 232–237.
- Hensch, T. (2005). Critical period plasticity in local cortical circuits. *Nat. Rev. Neurosci.* 6, 877–888.
- Herman, M. A., and Jahr, C. E. (2007). Extracellular glutamate concentration in hippocampal slice. *J. Neurosci.* 27, 9736–9741.
- Hirase, H., Qian, L., Barthó, P., and Buzsáki, G. (2004). Calcium dynamics of cortical astrocytic networks *in vivo*. *PLoS Biol.* 2:E96. doi: 10.1371/journal.pbio.0020096
- Houades, V., Koulakoff, A., Ezan, P., Seif, I., and Giaume, C. (2008). Gap junction-mediated astrocytic networks in the mouse barrel cortex. *J. Neurosci.* 28, 5207–5217.
- Iacobas, D. A., Suadicani, S. O., Spray, D. C., and Scemes, E. (2006). A stochastic two-dimensional model of intercellular Ca^{2+} wave spread in glia. *Biophys. J.* 90, 24–41.
- Iino, M. (1990). Biphasic Ca^{2+} -dependence of inositol 1,4,5-trisphosphate-induced Ca^{2+} release in smooth muscle cells of the guinea pig *Taenia caeci*. *J. Gen. Physiol.* 95, 1103–1112.
- Irvine, R. F., Lloyd-Burton, S. M., Yu, J. C. H., Letcher, A. J., and Schell, M. J. (2006). The regulation and function of inositol 1,4,5-trisphosphate 3-kinase. *Adv. Enzyme Regul.* 46, 314–323.
- Jaiswal, J., Fix, M., Takano, T., Nedergaard, M., and Simon, S. (2007). Resolving vesicle fusion from lysis to monitor calcium-triggered lysosomal exocytosis in astrocytes. *Proc. Natl. Acad. Sci. U.S.A.* 104, 14151–14156.
- Jia, H., Rochefort, N., Chen, X., and Konnerth, A. (2010). Dendritic organization of sensory input to cortical neurons *in vivo*. *Nature* 464, 1307–1312.
- Jourdain, P., Bergersen, L. H., Bhaukaurally, K., Bezzi, P., Santello, M., Domercq, M., et al. (2007). Glutamate exocytosis from astrocytes controls synaptic strength. *Nat. Neurosci.* 10, 331–339.
- Kaftan, E. J., Erlich, B. E., and Watras, J. (1997). Inositol 1,4,5-trisphosphate (InsP₃) and calcium interact to increase the dynamic range of InsP₃ receptor-dependent calcium signaling. *J. Gen. Physiol.* 110, 529–538.
- Kang, J., Jiang, L., Goldman, S. A., and Nedergaard, M. (1998). Astrocyte-mediated potentiation of inhibitory synaptic transmission. *Nat. Neurosci.* 1, 683–692.
- Kang, M., and Othmer, H. (2009). Spatiotemporal characteristics of calcium dynamics in astrocytes. *Chaos* 19, 037116.
- Kang, N., Xu, J., Xu, Q., Nedergaard, M., and Kang, J. (2005). Astrocytic glutamate release-induced transient depolarization and epileptiform discharges in hippocampal CA1 pyramidal neurons. *J. Neurophysiol.* 94, 4121–4130.
- Keizer, J., Li, Y., Stojilkovic, S., and Rinzler, J. (1995). InsP₃-induced Ca^{2+} excitability of the endoplasmic reticulum. *Mol. Biol. Cell* 6, 945–951.
- Kozlov, A. S., Angulo, M. C., Audinat, E., and Charpak, S. (2006). Target cell-specific modulation of neuronal activity by astrocytes. *Proc. Natl. Acad. Sci. U.S.A.* 103, 10058–10063.
- Kreft, M., Stenovec, M., Rupnik, M., Grilc, S., Kržan, M., Potokar, M., et al. (2004). Properties of Ca^{2+} -dependent exocytosis in cultured astrocytes. *Glia* 46, 437–445.
- Kuga, N., Sasaki, T., Takahara, Y., Matsuki, N., and Ikegaya, Y. (2011). Large-scale calcium waves traveling through astrocytic networks *in vivo*. *J. Neurosci.* 31, 2607–2614.
- Kulik, A., Haentzsch, A., Lüchermann, M., Reichelt, W., and Ballanyi, K. (1999). Neuron-glia signaling via α_1 adrenoreceptor-mediated Ca^{2+} release in Bergmann glial cells *in situ*. *J. Neurosci.* 19, 8401–8408.
- Lallouette, J., and Berry, H. (2012). “Topology drives calcium wave propagation in 3D astrocyte networks,” in *European Conference on Complex Systems* 2012, Bruxelles.
- Larsson, M., Sawada, K., Morland, C., Hiasa, M., Ormel, L., Moriyama, Y., et al. (2011). Functional and anatomical identification of a vesicular transporter mediating neuronal ATP release. *Cereb. Cortex* 22, 1203–1214.
- Lavialle, M., Aumann, G., Anlauf, E., Pröls, F., Arpin, M., and Derouiche, A. (2011). Structural plasticity of perisynaptic astrocyte processes involves ezrin and metabotropic glutamate receptors. *Proc. Natl. Acad. Sci. U.S.A.* 108, 12915–12919.
- Lee, M., McGeer, E., and McGeer, P. (2011). Mechanisms of GABA release from human astrocytes. *Glia* 59, 1600–1611.
- Lee, S., Yoon, B., Berglund, K., Oh, S., Park, H., Shin, H., et al. (2010). Channel-mediated tonic GABA release from glia. *Science* 330, 790–796.
- Le Meur, K., Mendizabal-Zubiaga, J., Grandes, P., and Audinat, E. (2012). GABA release by hippocampal astrocytes. *Front. Comput. Neurosci.* 6:59. doi: 10.3389/fncom.2012.00059
- Li, D., Ropert, N., Koulakoff, A., Giaume, C., and Oheim, M. (2008). Lysosomes are the major vesicular compartment undergoing Ca^{2+} -regulated exocytosis from cortical astrocytes. *J. Neurosci.* 28, 7648–7658.
- Li, L., Bender, K., Drew, P., Jadhav, S., Sylwestrak, E., and Feldman, D. (2009). Endocannabinoid signaling is required for development and critical period plasticity of the whisker map in somatosensory cortex. *Neuron* 64, 537–549.
- Li, Y.-X., Rinzler, J., Keizer, J., and Stojilkovic, S. S. (1994). Calcium oscillations in pituitary gonadotrophs: comparison of experiment and theory. *Proc. Natl. Acad. Sci. U.S.A.* 91, 58–62.
- Liu, Q., Xu, Q., Arcuino, G., Kang, J., and Nedergaard, M. (2004a). Astrocyte-mediated activation of neuronal kainate receptors. *Proc. Natl. Acad. Sci. U.S.A.* 101, 3172–3177.
- Liu, Q.-S., Xu, Q., Kang, J., and Nedergaard, M. (2004b). Astrocyte activation of presynaptic metabotropic glutamate receptors modulates hippocampal inhibitory synaptic transmission. *Neuron Glia Biol.* 1, 307–316.
- Lovatt, D., Xu, Q., Liu, W., Takano, T., Smith, N. A., Schnerrmann, J., et al. (2012). Neuronal adenosine release, and not astrocytic ATP release, mediates feedback inhibition of excitatory activity. *Proc. Natl. Acad. Sci. U.S.A.* 109, 6265–6270.
- Mak, D., McBride, S., and Foskett, J. (2003). Spontaneous channel activity of the inositol 1,4,5-trisphosphate (InsP₃) receptor (InsP₃R). Application of allosteric modeling to calcium and InsP₃ regulation of InsP₃R single-channel gating. *J. Gen. Physiol.* 122, 583–603.
- Malarkey, E., Ni, Y., and Parpura, V. (2008). Ca^{2+} entry through TRPC1 channels contributes to intracellular Ca^{2+} dynamics and consequent glutamate release from rat astrocytes. *Glia* 56, 821–835.
- Malarkey, E., and Parpura, V. (2011). Temporal characteristics of vesicular fusion in astrocytes: examination of synaptobrevin 2-laden vesicles at single vesicle resolution. *J. Physiol.* 589, 4271–4300.
- Marchaland, J., Cali, C., Voglmaier, S. M., Li, H., Regazzi, R., Edwards, R. H., et al. (2008). Fast subplasma membrane Ca^{2+} transients control exo-endocytosis of synaptic-like microvesicles in astrocytes. *J. Neurosci.* 28, 9122–9132.
- Matyash, V., Filippov, V., Mohrhagen, K., and Kettenmann, H. (2001). Nitric oxide signals parallel fiber activity to Bergmann glial cells in the mouse cerebellar slice. *Mol. Cell. Neurosci.* 18, 664–670. Abstract.
- Matyash, V., and Kettenmann, H. (2010). Heterogeneity in astrocyte morphology and physiology. *Brain Res. Rev.* 63, 2–10.
- Min, R., and Nevian, T. (2012). Astrocyte signaling controls spike timing-dependent depression at neocortical synapses. *Nat. Neurosci.* 15, 746–753.
- Mishra, J., and Bhalla, U. S. (2002). Simulations of inositol phosphate metabolism and its interaction with InsP₃-mediated calcium release. *Biophys. J.* 83, 1298–1316.
- Montana, V., Malarkey, E. B., Verderio, C., Matteoli, M., and Parpura, V. (2006). Vesicular transmitter release from astrocytes. *Glia* 54, 700–715.
- Montana, V., Ni, Y., Sunjara, V., Hua, X., and Parpura, V. (2004). Vesicular glutamate transporter-dependent glutamate release from astrocytes. *J. Neurosci.* 24, 2633–2642.
- Mulligan, S., and MacVicar, B. (2004). Calcium transients in astrocyte endfeet cause cerebrovascular constrictions. *Nature* 431, 195–199.
- Muyderman, H., Ångehagen, M., Sandberg, M., Björklund, U., Olsson, T., Hansson, E., et al. (2001). α_1 -Adrenergic modulation of metabotropic glutamate receptor-induced calcium oscillations and glutamate release in astrocytes. *J. Biol. Chem.* 276, 46504–46514.
- Navarrete, M., and Araque, A. (2008). Endocannabinoids mediate neuron-astrocyte communication. *Neuron* 57, 883–893.
- Navarrete, M., and Araque, A. (2010). Endocannabinoids potentiate synaptic transmission through stimulation of astrocytes. *Neuron* 68, 113–126.

- Navarrete, M., and Araque, A. (2011). Basal synaptic transmission: astrocytes rule! *Cell* 146, 675–677.
- Navarrete, M., Perea, G., de Sevilla, D., Gómez-Gonzalo, M., Núñez, A., Martn, E., et al. (2012a). Astrocytes mediate *in vivo* cholinergic-induced synaptic plasticity. *PLoS Biol.* 10:e1001259. doi: 10.1371/journal.pbio.1001259
- Navarrete, M., Perea, G., Maglio, L., Pastor, J., García de Sola, R., and Araque, A. (2012b). Astrocyte calcium signal and gliotransmission in human brain tissue. *Cereb. Cortex*. doi: 10.1093/cercor/bhs122. [Epub ahead of print].
- Nett, W. J., Oloff, S. H., and McCarthy, K. D. (2002). Hippocampal astrocytes *in situ* exhibit calcium oscillations that occur independent of neuronal activity. *J. Neurophysiol.* 87, 528–537.
- Newman, E. (2005). Calcium increases in retinal glial cells evoked by light-induced neuronal activity. *J. Neurosci.* 25, 5502–5510.
- Newman, E. A. (2003). Glial cell inhibition of neurons by release of ATP. *J. Neurosci.* 23, 1659–1666.
- Newman, E. A., and Zahs, K. R. (1997). Calcium waves in retinal glial cells. *Science* 275, 844–847.
- Ni, Y., Malarkey, E. B., and Parpura, V. (2007). Vesicular release of glutamate mediates bidirectional signaling between astrocytes and neurons. *J. Neurochem.* 103, 1273–1284.
- Nimmerjahn, A. (2009). Astrocytes going live: advances and challenges. *J. Physiol.* 587, 1639–1647.
- Nimmerjahn, A., Kirchhoff, F., Kerr, J. N. D., and Helmchen, F. (2004). Sulforhodamine 101 as a specific marker of astroglia in the neocortex *in vivo*. *Nat. Methods* 1, 31–37.
- Okubo, Y., Sekiya, H., Namiki, S., Sakamoto, H., Iinuma, S., Yamasaki, M., et al. (2010). Imaging extrasynaptic glutamate dynamics in the brain. *Proc. Natl. Acad. Sci. U.S.A.* 107, 6526–6531.
- Oliet, S. H. R., Piet, R., and Poulain, D. A. (2001). Control of glutamate clearance and synaptic efficacy by glial coverage of neurons. *Science* 292, 923–926.
- Panati, A., Theodosis, D. T., Mothet, J. P., Touquet, B., Pollegioni, L., Poulain, D. A., et al. (2006). Gliaderived D-serine controls NMDA receptor activity and synaptic memory. *Cell* 125, 775–784.
- Panati, A., Vallée, J., Haber, M., Murai, K., Lacaille, J., and Robitaille, R. (2011). Astrocytes are endogenous regulators of basal transmission at central synapses. *Cell* 146, 785–798.
- Pangršić, T., Potokar, M., Stenovec, M., Kreft, M., Fabbretti, E., Nistri, A., et al. (2007). Exocytotic release of ATP from cultured astrocytes. *J. Biol. Chem.* 282, 28749–28758.
- Pankratov, Y., Lalo, U., Verkhratsky, A., and North, R. (2007). Quantal release of ATP in mouse cortex. *J. Gen. Physiol.* 129, 257–265.
- Pannasch, U., Vargová, L., Reingruber, J., Ezan, P., Holcman, D., Giaume, C., et al. (2011). Astroglial networks scale synaptic activity and plasticity. *Proc. Natl. Acad. Sci. U.S.A.* 108, 8467–8472.
- Papouin, T., Ladépêche, L., Ruel, J., Sacchi, S., Labasque, M., Hanini, M., et al. (2012). Synaptic and extrasynaptic NMDA receptors are gated by different endogenous coagonists. *Cell* 150, 633–646.
- Parpura, V., Fang, Y., Basarsky, T., Jahn, R., and Haydon, P. G. (1995). Expression of synaptobrevin II, cellubrevin and syntaxin but not SNAP-25 in cultured astrocytes. *FEBS Lett.* 377, 489–492.
- Parpura, V., Grubišić, V., and Verkhratsky, A. (2011). Ca²⁺ sources for the exocytotic release of glutamate from astrocytes. *Biochim. Biophys. Acta* 1813, 984–991.
- Parpura, V., and Haydon, P. G. (2000). Physiological astrocytic calcium levels stimulate glutamate release to modulate adjacent neurons. *Proc. Natl. Acad. Sci. U.S.A.* 97, 8629–8634.
- Parpura, V., and Zorec, R. (2010). Gliotransmission: exocytotic release from astrocytes. *Brain Res. Rev.* 63, 83–92.
- Parri, H. R., Gould, T. M., and Crunelli, V. (2001). Spontaneous astrocytic Ca²⁺ oscillations *in situ* drive NMDAR-mediated neuronal excitation. *Nat. Neurosci.* 4, 803–812.
- Pascual, O., Casper, K. B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J. Y., et al. (2005). Astrocytic purinergic signaling coordinates synaptic networks. *Science* 310, 113–116.
- Pasti, L., Volterra, A., Pozzan, T., and Carmignoto, G. (1997). Intracellular calcium oscillations in astrocytes: a highly plastic, bidirectional form of communication between neurons and astrocytes *in situ*. *J. Neurosci.* 17, 7817–7830.
- Pasti, L., Zonta, M., Pozzan, T., Vicini, S., and Carmignoto, G. (2001). Cytosolic calcium oscillations in astrocytes may regulate exocytotic release of glutamate. *J. Neurosci.* 21, 477–484.
- Perea, G., and Araque, A. (2005). Properties of synaptically evoked astrocyte calcium signal reveal synaptic information processing by astrocyte. *J. Neurosci.* 25, 2192–2203.
- Perea, G., and Araque, A. (2007). Astrocytes potentiate transmitter release at single hippocampal synapses. *Science* 317, 1083–1086.
- Perea, G., Navarrete, M., and Araque, A. (2009). Tripartite synapse: astrocytes process and control synaptic information. *Trends Neurosci.* 32, 421–431.
- Petravicz, J., Fiacco, T., and McCarthy, K. (2008). Loss of IP₃ receptor-dependent Ca²⁺ increases in hippocampal astrocytes does not affect baseline CA1 pyramidal neuron synaptic activity. *J. Neurosci.* 28, 4967–4973.
- Piet, R., and Jahr, C. (2007). Glutamatergic and purinergic receptor-mediated calcium transients in Bergmann glial cells. *J. Neurosci.* 27, 4027–4035.
- Piet, R., Vargová, L., Syková, E., Poulain, D., and Oliet, S. (2004). Physiological contribution of the astrocytic environment of neurons to intersynaptic crosstalk. *Proc. Natl. Acad. Sci. U.S.A.* 101, 2151–2155.
- Pinheiro, P. S., and Mulle, C. (2008). Presynaptic glutamate receptors: physiological functions and mechanisms of action. *Nat. Rev.* 9, 423–436.
- Pirttimäki, T., Hall, S., and Parri, H. (2011). Sustained neuronal activity generated by glial plasticity. *J. Neurosci.* 31, 7637–7647.
- Pivneva, T., Haas, B., Reyes-Haro, D., Laube, G., Veh, R., Nolte, C., et al. (2008). Store-operated Ca²⁺ entry in astrocytes: different spatial arrangement of endoplasmic reticulum explains functional diversity *in vitro* and *in situ*. *Cell Calcium* 43, 591–601.
- Porter, J. T., and McCarthy, K. D. (1996). Hippocampal astrocytes *in situ* respond to glutamate released from synaptic terminals. *J. Neurosci.* 16, 5073–5081.
- Poskanzer, K. E., and Yuste, R. (2011). Astrocytic regulation of cortical UP states. *Proc. Natl. Acad. Sci. U.S.A.* 108, 18453–18458.
- Pryazhnikov, E., and Khiroug, L. (2008). Sub-micromolar increase in [Ca²⁺]_i triggers delayed exocytosis of ATP in cultured astrocytes. *Glia* 56, 38–49.
- Raghavachari, S., and Lisman, J. E. (2004). Properties of quantal transmission at CA1 synapses. *J. Neurophysiol.* 92, 2456–2467.
- Ramos-Franco, J., Bare, D., Caenepeel, S., Nani, A., Fill, M., and Mignery, G. (2000). Single-channel function of recombinant type 2 inositol 1,4,5-trisphosphate receptor. *Biophys. J.* 79, 1388–1399.
- Rebecchi, M. J., and Pentylala, S. N. (2000). Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiol. Rev.* 80, 1291–1335.
- Rebola, N., Rodrigues, R., Lopes, L., Richardson, P., Oliveira, C., and Cunha, R. (2005). Adenosine A₁ and A_{2A} receptors are co-expressed in pyramidal neurons and co-localized in glutamatergic nerve terminals of the rat hippocampus. *Neuroscience* 133, 79–83.
- Regehr, W. F., Delaney, K. R., and Tank, D. W. (1994). The role of presynaptic calcium in short-term enhancement at the hippocampal mossy fiber synapse. *J. Neurosci.* 14, 523–537.
- Rieger, A., Deitmer, J., and Lohr, C. (2007). Axon-glia communication evokes calcium signaling in olfactory ensheathing cells of the developing olfactory bulb. *Glia* 55, 352–359.
- Robitaille, R. (1998). Modulation of synaptic efficacy and synaptic depression by glial cells at the frog neuromuscular junction. *Neuron* 21, 847–855.
- Rodríguez-Moreno, A., and Paulsen, O. (2008). Spike timing-dependent long-term depression requires presynaptic nmda receptors. *Nat. Neurosci.* 11, 744–745.
- Rossi, D. (2012). Astrocytes join the plasticity party. *Nat. Neurosci.* 15, 649–651.
- Rouach, N., Koulakoff, A., and Giaume, C. (2004). Neurons set the tone of gap junctional communication in astrocytic networks. *Neurochem. Int.* 45, 265–272.
- Rousse, I., St-Amour, A., Darabid, H., and Robitaille, R. (2010). Synapse-glia interactions are governed by synaptic and intrinsic glial properties. *Neuroscience* 167, 621–632.
- Roux, L., Benchenane, K., Rothstein, J., Bonvento, G., and Giaume, C. (2011). Plasticity of astroglial networks in olfactory glomeruli. *Proc. Natl. Acad. Sci. U.S.A.* 108, 18442–18446.
- Santello, M., Bezzi, P., and Volterra, A. (2011). TNF α controls glutamatergic gliotransmission in the hippocampal dentate gyrus. *Neuron* 69, 988–1001.
- Santello, M., and Volterra, A. (2009). Synaptic modulation by astrocytes

- via Ca^{2+} -dependent glutamate release. *Neuroscience* 158, 253–259.
- Santello, M., and Volterra, A. (2010). Astrocytes as aide-mémoires. *Nature* 463, 169–170.
- Santello, M., and Volterra, A. (2012). TNF α in synaptic function: switching gears. *Trends Neurosci.* 35, 638–647.
- Sasaki, T., Matsuki, N., and Ikegaya, Y. (2011). Action-potential modulation during axonal conduction. *Science* 331, 599–601.
- Sawada, K., Echigo, N., Juge, N., Miyaji, T., Otsuka, M., Omote, H., et al. (2008). Identification of a vesicular nucleotide transporter. *Proc. Natl. Acad. Sci. U.S.A.* 105, 5683–5686.
- Sawyer, C., and Clifton, D. (1980). Aminergic innervation of the hypothalamus. *Fed. Proc.* 39, 2889–2895.
- Scemes, E., and Giaume, C. (2006). Astrocyte calcium waves: what they are and what they do. *Glia* 54, 716–725.
- Schipke, C., Haas, B., and Kettenmann, H. (2008). Astrocytes discriminate and selectively respond to the activity of a subpopulation of neurons within the barrel cortex. *Cereb. Cortex* 18, 2450–2459.
- Schmitt, L., Sims, R., Dale, N., and Haydon, P. (2012). Wakefulness affects synaptic and network activity by increasing extracellular astrocyte-derived adenosine. *J. Neurosci.* 32, 4417–4425.
- Schubert, V., Bouvier, D., and Volterra, A. (2011). SNARE protein expression in synaptic terminals and astrocytes in the adult hippocampus: a comparative analysis. *Glia* 59, 1472–1488.
- Schummers, J., Yu, H., and Sur, M. (2008). Tuned responses of astrocytes and their influence on hemodynamic signals in the visual cortex. *Science* 320, 1638.
- Sejnowski, T. J., and Paulsen, O. (2006). Network oscillations: emerging computational principles. *J. Neurosci.* 26, 1673–1676.
- Serrano, A., Haddjeri, N., Lacaille, J., and Robitaille, R. (2006). GABAergic network activation of glial cells underlies heterosynaptic depression. *J. Neurosci.* 26, 5370–5382.
- Shadlen, M., and Newsome, W. (1994). Noise, neural codes and cortical organization. *Curr. Opin. Neurobiol.* 4, 569–579.
- Shigemoto, R., Kinoshita, A., Wada, E., Nomura, S., Ohishi, H., Takada, M., et al. (1997). Differential presynaptic localization of metabotropic glutamate receptor subtypes in the rat hippocampus. *J. Neurosci.* 17, 7503–7522.
- Shigetomi, E., Tong, X., Kwan, K., Corey, D., and Khakh, B. (2011). TRPA1 channels regulate astrocyte resting calcium levels and inhibitory synapse efficacy via GAT-3. *Nat. Neurosci.* 15, 70–80.
- Shinozaki, T., Michikawa, T., Enomoto, M., Goto, J., Iwai, M., Matsuura, T., et al. (2011). Mechanistic basis of bell-shaped dependence of inositol 1 4, 5-trisphosphate receptor gating on cytosolic calcium. *Proc. Natl. Acad. Sci. U.S.A.* 108, 15486–15491.
- Shu, Y., Hasenstaub, A., and McCormick, D. (2003). Turning on and off recurrent balanced cortical activity. *Nature* 423, 288–293.
- Sims, C. E., and Allbritton, N. L. (1998). Metabolism of inositol 1 4, 5-trisphosphate and inositol 1 3 4, 5-tetrakisphosphate by the oocytes of *Xenopus laevis*. *J. Biol. Chem.* 273, 4052–4058.
- Sjöström, P., Turrigiano, G., and Nelson, S. (2003). Neocortical LTD via coincident activation of presynaptic NMDA and cannabinoid receptors. *Neuron* 39, 641–654.
- Sneyd, J., Charles, A. C., and Sanderson, M. J. (1994). A model for the propagation of intracellular calcium waves. *Am. J. Physiol.* 266, C293–C302.
- Stellwagen, D., Beattie, E. C., Seo, J. Y., and Malenka, R. C. (2005). Differential regulation of AMPA receptor and GABA receptor trafficking by tumor necrosis factor- α . *J. Neurosci.* 25, 3219–3228.
- Stellwagen, D., and Malenka, R. (2006). Synaptic scaling mediated by glial TNF- α . *Nature* 440, 1054–1059.
- Stenovec, M., Kreft, S., Grilc, M., Pokotar, M., Kreft, M. E., Pangršič, T., et al. (2007). Ca^{2+} -dependent mobility of vesicles capturing anti-VGLUT1 antibodies. *Exp. Cell Res.* 313, 3809–3818.
- Sul, J.-Y., Orosz, G., Givens, R. S., and Haydon, P. G. (2004). Astrocytic connectivity in the hippocampus. *Neuron Glia Biol.* 1, 3–11.
- Svoboda, K., Denk, W., Kleinfeld, D., and Tank, D. (1997). *In vivo* dendritic calcium dynamics in neocortical pyramidal neurons. *Nature* 385, 161–165.
- Swadlow, H. (2002). Thalamocortical control of feed-forward inhibition in awake somatosensory ‘barrel’ cortex. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 357, 1717–1727.
- Syková, E., and Nicholson, C. (2008). Diffusion in brain extracellular space. *Physiol. Rev.* 88, 1277–1340.
- Takata, N., Mishima, T., Hisatsune, C., Nagai, T., Ebisui, E., Mikoshiba, K., et al. (2011). Astrocyte calcium signaling transforms cholinergic modulation to cortical plasticity *in vivo*. *J. Neurosci.* 31, 18155–18165.
- Thomas, D., Lipp, P., Tovey, S., Berridge, M., Li, W., Tsien, R., et al. (2000). Microscopic properties of elementary Ca^{2+} release sites in non-excitable cells. *Curr. Biol.* 10, 8–15.
- Tian, G.-F., Azmi, H., Takahiro, T., Xu, Q., Peng, W., Lin, J., et al. (2005). An astrocytic basis of epilepsy. *Nat. Med.* 11, 973–981.
- Tian, G. F., Takano, T., Lin, J. H.-C., Wang, X., Bekar, L., and Nedergaard, M. (2006). Imaging of cortical astrocytes using 2-photon laser scanning microscopy in the intact mouse brain. *Adv. Drug Deliv. Rev.* 58, 773–787.
- Todd, K. J., Darabid, H., and Robitaille, R. (2010). Perisynaptic glia discriminate patterns of motor nerve activity and influence plasticity at the neuromuscular junction. *J. Neurosci.* 30, 11870–11882.
- Torres, A., Wang, F., Xu, Q., Fujita, T., Dobrowolski, R., Willecke, K., et al. (2012). Extracellular Ca^{2+} acts as a mediator of communication from neurons to glia. *Sci. Signal.* 5, ra8.
- Troyer, T., and Miller, K. (1997). Physiological gain leads to high ISI variability in a simple model of a cortical regular spiking cell. *Neural Computat.* 9, 971–983.
- Ullian, E., Christopherson, K., and Barres, B. (2004). Role for glia in synaptogenesis. *Glia* 47, 209–216.
- van den Pol, A. N., Romano, C., and Ghosh, P. (1995). Metabotropic glutamate receptor mGluR5 subcellular distribution and developmental expression in hypothalamus. *J. Comp. Neurol.* 362, 134–150.
- Ventura, R., and Harris, K. M. (1999). Three-dimensional relationships between hippocampal synapses and astrocytes. *J. Neurosci.* 19, 6897–6906.
- Verkhatsky, A., Rodriguez, R. R., and Parpura, V. (2012). Calcium signalling in astroglia. *Mol. Cell. Endocrinol.* 353, 45–56.
- Vogels, T., and Abbott, L. (2009). Gating multiple signals through detailed balance of excitation and inhibition in spiking networks. *Nat. Neurosci.* 12, 483–491.
- Volman, V., Bazhenov, M., and Sejnowski, T. J. (2011). “A computational model of neuronal and glial homeostatic synaptic plasticity in posttraumatic epileptogenesis.” in *Neuroscience Meeting Planner*, Number 250.11/V5. Society for Neuroscience. [Online].
- Volman, V., Ben-Jacob, E., and Levine, H. (2007). The astrocyte as a gatekeeper of synaptic information transfer. *Neural Comput.* 19, 303–326.
- Volterra, A., and Meldolesi, J. (2005). Astrocytes, from brain glue to communication elements: the revolution continues. *Nat. Rev. Neurosci.* 6, 626–640.
- Wang, X., Lou, N., Xu, Q., Tian, G.-F., Peng, W. G., Han, X., et al. (2006). Astrocytic Ca^{2+} signaling evoked by sensory stimulation *in vivo*. *Nat. Neurosci.* 9, 816–823.
- Waters, J., Larkum, M., Sakmann, B., and Helmchen, F. (2003). Supralinear Ca^{2+} influx into dendritic tufts of layer 2/3 neocortical pyramidal neurons *in vitro* and *in vivo*. *J. Neurosci.* 23, 8558–8567.
- Watrás, J., Bezprozvanny, I., and Ehrlich, B. E. (1991). Inositol 1 4, 5-trisphosphate-gated channels in cerebellum: presence of multiple conductance states. *J. Neurosci.* 11, 3239–3245.
- Winship, I., Plaa, N., and Murphy, T. (2007). Rapid astrocyte calcium signals correlate with neuronal activity and onset of the hemodynamic response *in vivo*. *J. Neurosci.* 27, 6268–6272.
- Yang, Y., Ge, W., Chen, Y., Zhang, Z., Shen, W., Wu, C., et al. (2003). Contribution of astrocytes to hippocampal long-term potentiation through release of D-serine. *Proc. Natl. Acad. Sci. U.S.A.* 100, 15194–15199.
- Zhang, B.-X., Zhao, H., and Muallem, S. (1993). Calcium dependent kinase and phosphatase control inositol-1 4, 5-trisphosphate-mediated calcium release: modification by agonist stimulation. *J. Biol. Chem.* 268, 10997–11001.
- Zhang, J.-M., Wang, H.-K., Ye, C.-Q., Ge, W., Chen, Y., Jiang, Z.-L., et al. (2003). ATP released by astrocytes mediates glutamatergic activity-dependent heterosynaptic suppression. *Neuron* 40, 971–982.
- Zhang, Q., Fukuda, M., Van Bockstaele, E., Pascual, O., and Haydon, P. G. (2004a). Synaptotagmin IV regulates glial glutamate release. *Proc. Natl. Acad. Sci. U.S.A.* 101, 9441–9446.
- Zhang, Q., Pangršič, T., Kreft, M., Kržan, M., Li, N., Sul, J.-Y., et al. (2004b). Fusion-related release of glutamate from astrocytes. *J. Biol. Chem.* 279, 12724–12733.

- Zhang, Y., and Barres, B. (2010). Astrocyte heterogeneity: an underappreciated topic in neurobiology. *Curr. Opin. Neurobiol.* 20, 588–594.
- Zhang, Z., Chen, G., Zhou, W., Song, A., Xu, T., Luo, Q., et al. (2007). Regulated ATP release from astrocytes through lysosome exocytosis. *Nat. Cell Biol.* 9, 945–953.
- Zonta, M., and Carmignoto, G. (2002). Calcium oscillations encoding neuron-to-astrocyte communication. *J. Physiol. Paris* 96, 193–198.
- Zorec, R., Araque, A., Carmignoto, G., Haydon, P., Verkhratsky, A., and Parpura, V. (2012). Astroglial excitability and gliotransmission: an appraisal of Ca^{2+} as a signaling route. *ASN Neuro* 4, e00080.
- Zucker, R. S., and Regehr, W. G. (2002). Short-term synaptic plasticity. *Annu. Rev. Physiol.* 64, 355–405.
- Zur Nieden, R., and Deitmer, J. W. (2006). The role of metabotropic glutamate receptors for the generation of calcium oscillations in rat hippocampal astrocytes *in situ*. *Cereb. Cortex* 16, 676–687.
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Received: 23 April 2012; paper pending published: 28 May 2012; accepted: 06 December 2012; published online: 21 December 2012.
- Citation: De Pittà M, Volman V, Berry H, Parpura V, Volterra A, and Ben-Jacob E (2012) Computational quest for understanding the role of astrocyte signaling in synaptic transmission and plasticity. *Front. Comput. Neurosci.* 6:98. doi: 10.3389/fncom.2012.00098
- Copyright © 2012 De Pittà, Volman, Berry, Parpura, Volterra, and Ben-Jacob. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and subject to any copyright notices concerning any third-party graphics etc.